

EVALUATION OF THE IMMUNE RESPONSE TO POLIO VACCINE IN MALNOURISHED CHILDREN IN SANA'A CITY

Abstract

This study was made to evaluate the immune response to polio virus vaccine among PEM children by measuring the level of circulating Immunoglobulin G (IgG) antibodies against polio virus (IgG-PV) after immunization with the primary series of POV, and determining the coverage rate of universal childhood vaccine for polio virus. A cross-sectional laboratory study was conducted in Department of Medical Microbiology and Clinical Immunology, Faculty of Medicine and Health Sciences, and Al-Sabeen University Hospital, Sana'a University. A total of 279 PEM children were selected and investigated for universal childhood vaccination coverage rate for polio vaccine. Blood samples were collected from all, then tested for levels of IgG-PV by ELISA method. For assessment IgG-PV levels more than 10 units/ml were considered protected against polio virus infection.

The coverage rate of polio virus vaccine for first year vaccine was 96.8%; and 91.1% of vaccinated PEM children responded to the vaccine with mean level of 46.2 U/ml. A statistically significant difference was observed with respect to sero-protective IgG-PV between males and females (85.7% and 94.1% respectively, $p=0.002$); and older children (>37 months) (97.7%).

We conclude that a small proportion of malnourished vaccinated children with a normal immune status were not serologically immune to polio virus infection, and remain to be reconsidered for either revaccination or booster doses due to lack of or inadequate response. PEM group gave slightly reduced response to OPV hence there is need to give this group IPV (injectable polio vaccine) along with OPV and different micro-nutrition deficiencies like Zinc and iron.

Keywords: children, malnutrition, PEM, polio vaccine, POV, Sana'a city, Yemen

Introduction

Globally Malnutrition is the most important cause for failure of all vaccination.^{1,2} Consistent with Independent Monitoring Board of the Global Polio Eradication Initiative, Tenth Report 2014 on 28th of October 2014, globally 257 polio cases were reported in three polio endemic countries, of which 220, wild polio cases were reported from Pakistan.^{2,3,4} About 165 million children are universally, chronically malnourished. This preventable state has affected one in every four children. Malnutrition cause 2.3 million children deaths per year, an average of one death per every 15 seconds.⁵ According to WHO majority of expected malnourished children live in developing countries. Less than 30% of them are under the age of five years and half of them are suffering from PEM.^{1,2} The report of MHP Yemen, and UNICEF shows the PEM is very high in Yemen among children (50%) this Malnutrition might be one of the main difficulty for polio eradication among under five years old children in Yemen.⁶ In its 41st meeting in 1988, World Health Assembly approved a declaration according to which "OPV is exclusively used worldwide for polio eradication for some reasons. It is cheaper than Intramuscular polio Vaccine (IPV) and can be easily orally administered".² However, worldwide, the

oral vaccine doesn't effectively develop immunity against polio, particularly in malnourished children.⁷ Some data from WHO has also highlighted that malnutrition hinders the battle against polio because OPV produces four times less immunity in malnourished children as compared to well-nourished. Since 50% of under-five children are malnourished in Yemen, it is very difficult to eradicate polio from Yemen.⁶ Therefore, our priority should be focused on preventing malnutrition in this group of population.

This study was made to evaluate the immune response to polio virus vaccine among PEM children by measuring the level of circulating Immunoglobulin G (IgG) antibodies against polio virus (IgG-PV) after immunization with the primary series of POV, and determining the coverage rate of universal childhood vaccine for polio virus.

Subjects and Methods

Ethical Consideration

Ethical clearance for the study was taken from the Faculty of Medicine and Health Sciences Research Review Committee. A written permission was also taken from the head of the pediatric department of the Al-Sabian Hospital, Sana'a city, Yemen. Informed Consent was taken from the parents and guardians of all selected children before the questionnaire was filled in and before the blood samples were collected.

Survey procedure and Laboratory Analysis

Demographic data and other relevant information of each participant were obtained through a questionnaire. Then, about 4 ml of blood was aseptically drawn by venipuncture after swabbing the area of interest for sample collection with alcohol. The serum was then separated from the blood by allowing clotting and centrifuging. Finally, the serum samples were labeled and stored at -20°C until processed. The polio IgG antibody ELISA test kit, manufactured and described by DEMEDITEC Diagnostic GmbH Germany, was used for the detection of specific IgG antibodies against polio in the serum of children with strict adherence to the manufacturer's instruction manual.

Study setting

The study was conducted at the pediatric department of the Al-Sabian Hospital, Sana'a city, Yemen. The Hospital was selected because it is the only teaching hospitals in Sana'a city provides health services to malnutrition children.

Study Population

Children between the age of 6 months to 59 months who attended the pediatric patient Department (PD) with PEM. These subjects of the study were take on from 17th of March, 2017 to 17th of September 2017 as the study population. It was ensured that they must have received first year oral polio vaccine doses and fulfill the eligibility criteria of study. A total of 279 blood samples were collected.

Sample size

The sample size was calculated by using sample size determination in health studies by WHO with 95% confidence interval and 5% level of significance with acceptable margin of error equal to 6% with PEM population in Yemen at least equal to 500 000 children . The proportions of antibodies production among malnourished subjects ("Evaluation of the response to vaccination against poliomyelitis and measles in malnourished children"⁸

was 59.5% in malnutrition children. The sample size was calculated to be 257 subjects . In such a small sample size, the chance of type 1 and type 2 errors are increased. To reduce these errors in the study, we increased the sample size to 279 subjects.

Result

The study included 270 malnourished children (126 males and 153 females). Most of our children patients were in age group 13-24 months in which they count 36.9%, followed by age group 3-12 months, in which the rate was 31.2% (table 1).

The non-protective of polio-antibodies IgG rate in our children patients was 8.6%, while the protective rate of polio-antibodies IgG in them was 91.4% (table 2). There was 8.6% of malnourished children had 2-9.9IU/ml Polio-virus IgG antibodies (susceptibility to polio infection), 28.3% of them had 11-30IU/ml (deprived protective level, and need urgent booster vaccine), 42.7% of malnourished children had 31-50IU/ml (needy protective level which might faint in short time, and need booster vaccine), 10.4% of malnourished children had 51-70IU/ml Polio-virus IgG antibodies (good health protective level) and 10% of malnourished children had ≥ 71 IU/ml polio-virus IgG antibodies (fabulous protective level) (table 3). The susceptibility rate to polio virus infection among males was 14.3%, higher than that of females 3.9%, with OR of susceptibility to poliovirus for male equal to 4.1 times than female ($p=0.002$). As regard to the age, the susceptibility rates were increase in younger age groups (table 4). Table 5 shows the associated odds ratio of susceptibility to polio virus infection with history of polio vaccine of malnourished children. There was high susceptibility rate to polio infection with children delivery in hospital (9.1%), and history of first month illness (9.5%), and in children had mixed feeding (33.3%), but no statistical significance was observed (table 6).

Discussion

Yemen introduced universal immunization against Polio virus infection for infants (POV) in early 70s of the last century, but feedback on the coverage rate of vaccinations and their efficacy in the community have been ignored for a long period. In addition, information on the prevalence and risk determinants of polio virus infection and on vaccination coverage rate and immune status against polio virus infection among children particularly malnourished children in Yemen has been inadequate or non-existent. Consequently, this study has been carried out to help our understanding of some of these questions. In addition, seroprevalence studies provide important data on act of immunization programs, susceptible groups and populations at-risk of future outbreaks. Identifying risk factors that affect seroconversion of the oral polio vaccine (OPV) will enable the polio eradication initiatives to increase seroprevalence. This study demonstrates the first hospital-based study aiming to determine the immune status of malnourished children in Sana'a city against polio virus. A total of 279 serum samples from malnourished children in Sana'a city were collected and analyzed in this study for polio-specific IgG antibodies, there was 91.4% of children had protective of polio-antibodies IgG against polio virus infection, while 8.6% of tested children had non protective level of polio-antibodies IgG against polio virus infection, with other words they are susceptible and under risk of polio virus infections. The high seroprevalence of 91.1% recorded in this study among malnourished children in Sana'a city indicates children have effectively responded to the vaccine being used in the ongoing polio eradication initiative in spite of malnourished. These findings are similar with United States study in which it was reported that 90% of healthy school age children, adolescent

and young adults had detectable antibodies to Poliovirus in spite of their nourished.⁹ However, present seroprevalence of 91.1% is higher than the findings of Egypt by El-Sayed *et al.*¹⁰, (37.8%) and Morocco by Yousuf *et al.*⁸ (21.4%), among malnourished children groups.

The current study seroprevalence of protective IgG antibodies for polio virus was slightly higher (94.1%) in females compared to males (85.7%), this result is different from study by Iiyasu *et al.*¹¹ in Nigeria, but similar to Donbraye *et al.*¹² study in Nigeria where females had higher antibody titers for all the three serotypes. Also our result is similar to Baba *et al.*¹³ in India and to Chinese study by Wang *et al.*¹⁴ also. Sex differential non-specific effects of vaccines are common in developing countries with negative non-specific effects (NSE) of inactivated vaccines more common in girls than boys^{16,17} in some countries, but with no differences in high income countries.¹⁷ Though the general pattern is both negative and positive NSE are stronger in females¹⁸, a randomized controlled trial in Guinea-Bissau negated the hypothesis that mortality rates in boys would be lower if they had not received OPV.¹⁹ The reasons why OPV uptake is slightly higher for females in our study area are unknown. Though factors such as age, gender and urban setting have no remedial solution from a public health perspective, this study indicates that response may be affected by factors that are not amenable to modification. This is important for the understanding of oral vaccine performance in low-income countries. When we considered age groups, there was trend toward increased rate of positive protective levels of IgG antibodies against polio virus in the current study in which the higher seropositivity rate was in the older age group (≥ 37 months age group) (97.7%) and the lowest protective rate was 87.4% in 6-12 months age group (table 4). This similar to findings in previous studies in Vellore district of Tamil Nadu - Sri Lanka in which lower rate of protective antibodies rate was found in children under one years of age.²⁰ This could be due to a more mature immune system or to receipt of OPV doses that are not recorded in the vaccination history taken for the child.

When we study the associated odds ratio of susceptibility to polio virus infection with history of polio vaccine among malnourished children in Sana'a city Yemen, there was increased rate of susceptibility for polio virus in our study among children whom not had national camping booster polio vaccine in which the rate of susceptibility was 14.6%, while the rate of susceptibility among children whom had national camping booster polio vaccine was 5.5% only (table 5). Our result is similar to previously reported studies in which increasing the number of OPV doses received by children, decreasing the susceptibility rate for polio virus; and increase the polio-seroconversion rates among children.^{8,11,21}

Conclusions

We conclude that a small proportion of malnourished vaccinated children with a normal immune status were not serologically immune to polio virus infection, and remain to be reconsidered for either revaccination or booster doses due to lack of or inadequate response; more information, however, needs to be gathered first on the occurrence of clinical/subclinical poliomyelitis cases (morbidity and mortality) and the overall morbidity and mortality following polio vaccination. Also the rates of vaccine coverage for the routine immunization schedule of childhood vaccines were good.

Acknowledgments:

Authors acknowledge the financial support of Sana'a University, Yemen.

Conflict of interest:

"No conflict of interest associated with this work".

References

- 1-WHO. "Polio vaccines: WHO position paper, March, 2016." . Wkly Epidemiol Rec 2016; **91** (12): 145–168.
- 2-WHO. "Poliomyelitis". World Health Organization. Archived from the original on 18 April 2017.
- 3-Caidi H, Bennis IF, Mouan N, El Aouad R. Evaluation of the response to vaccination against poliomyelitis and measles in malnourished children in Morocco. *East Mediterr Health J.* 2004;10(4-5):474–481.
- 4-World Health Organization: National Surveillance Bulletin Pakistan. 2011
- 5- Jack Dean. "165 million children malnourished worldwide". World Socialist Web Site Published by the International Committee of the Fourth International (ICFI), 1 June; 2013.
- 6-MHP (Ministry of Health and Population, Sana', Yemen). National Surveillance Health report of Yemen. 2017.
- 7-Sack DA, Qadri F, Svennerholm AM. Determinants of responses to oral vaccines in developing countries. *Ann Nestlé (Engl Ed)* 2008;66(2):71–79.
- 8-Yousuf A, Shah SA, Jaffery IA *et al.* Seroprevalence rate of Poliovirus antibodies among the Healthy and Protein Energy Malnutrition children. *Pak J Med Sci* 2015, 31(2):403-407.
- 9-Kelley PW, Petrucci BP, Stehr-Green P, Manson CJ. The susceptibility of young adult Americans to vaccine-preventable infections, natural sero-survey of US army recruits. *J Am Med Assoc.* 1991;266(19):2724–2729.
- 10-El-Sayed N, Al-Jorf S, Hennessey KA, Salama M, Watkins MA, Abdelwahab JA, *et al.* Survey of poliovirus antibodies during the final stage of polio eradication in Egypt. *Vaccine.* 2007;25(27):5062–5070.
- 11-Iliyasu Z., Nwaze E., Verma H., Mustapha A.O., Weldegebriel G., Gasasira A. Survey of poliovirus antibodies in Kano. Northern Nigeria. *Vaccine.* 2014;32(12):1414–1420.
- 12-Donbraye E., Adewumi M.O., Odaibo G.N., Bakarey A.S., Opaleye O.O., Olaleye D.O. Evaluation of immunity against poliovirus serotypes among children in Riverine areas of Delta State, Nigeria. *African J Clin Exp Microbiol.* 2011;12(2):72–75.
- 13-Baba M.M., Haruna B.A., Ogunmola O., Ambe J.P., Shidali N.N., Oderinde B. A survey for neutralizing antibodies to the three types of poliovirus among children in Maiduguri, Nigeria. *J Med Virol.* 2012;84(4):691–696.
- 14-Wang H., Cui H., Ding Z., Ba P., Zhu S., Wen N. Seroprevalence of anti-polio antibodies among children <15 years of age in border provinces in China. *Clin Vaccine Immunol.* 2013;20(7):1070–1075.
- 15-Aaby P., Jensen H., Gomes J., Fernandes M., Lisse I.M. The introduction of diphtheria-tetanus-pertussis vaccine and child mortality in rural Guinea-Bissau: an observational study. *Int J Epidemiol.* 2004;33(2):374–380.
- 16-Aaby P., Jensen H., Walraven G. Age-specific changes in the female-male mortality ratio related to the pattern of vaccinations: an observational study from rural Gambia. *Vaccine.* 2006;24(22):4701–4708.
- 17-Schurink-van't Klooster T.M., Knol M.J., de Melker H.E., van der Sande M.A. Gender-specific mortality in DTP-IPV- and MMR+/-MenC-eligible age groups to determine possible sex-differential effects of vaccination: an observational study. *BMC Infect Dis.* 2015;15:148.
- 18-Flanagan K.L., van Crevel R., Curtis N., Shann F., Levy O., Optimunize N. Heterologous ("nonspecific") and sex-differential effects of vaccines: epidemiology,

clinical trials, and emerging immunologic mechanisms. Clin Infect Dis. 2013;57(2):283–289.

19-Lund N., Andersen A., Hansen A.S., Jepsen F.S., Barbosa A., Biering-Sorensen S. The effect of oral polio vaccine at birth on infant mortality: a randomized trial. Clin Infect Dis. 2015;61(10):1504–1511.

20-Gamage D., Palihawadana P., Mach O., Weldon W.C., Oberste S.M., Sutter R.W. Achieving high seroprevalence against polioviruses in Sri Lanka-Results from a serological survey, 2014. J Epidemiol Glob Health. 2015;5(4 Suppl. 1):S67–S71.

21-Staat M.A., Stadler L.P., Donauer S., Trehan I., Rice M., Salisbury S. Serologic testing to verify the immune status of internationally adopted children against vaccine preventable diseases. Vaccine. 2010;28(50):7947–7955.

Table 1: The age and sex distribution of malnourished children, whom tested for immune status against polio virus, Sana'a city, Yemen

Age groups	Male		Female		Total	
	No	%	No	%	No	%
6 -12 months	42	33.3	45	29.4	87	31.2
13-24 months	48	38.1	55	35.9	103	36.9
25-36 months	18	14.3	27	17.6	45	16.1
≥ 37 months	18	14.3	26	17	44	15.8
Total	126	45.2	153	54.8	279	100

Table 2: The immune status of malnourished children against polio virus, Sana'a city, Yemen

Immune status	Number	percentage	<i>P V</i>
Non-protective level of polio-antibodies IgG (≤ 9.9 IU/ml)	24	8.6	< 0.05
Protective level of polio-antibodies IgG (≥ 10 IU/ml)	255	91.4	
Total	279	100	

Table 3: The interpretations of the poliomyelitis virus IgG antibodies level among malnourished children, Sana'a city, Yemen

Polio-virus IgG antibodies IU/ml	Number	Percentage	Interpretations'
2 - 9.9 IU/ml	24	8.6	Susceptible to polio infection
11- 30. IU/ml	79	28.3	deprived protective level, need urgent booster vaccine
31- 50 IU/ml	119	42.7	needy Protective level which might faint in short time, need booster vaccine
51 - 70 IU/ml	29	10.4	in good health protective level
≥ 71 IU/ml	28	10	fabulous protective level

Table 4: The frequency and associated odds ratio of susceptibility to polio virus infection with different sexes and age groups of malnourished children in Sana'a city Yemen.

Characters	Non-protective level N=24		OR	CI	χ^2	PV
	No	%				
Sex:						
Male n=126	18	14.3	4.1	1.6-10.6	9.4	0.002
Female n=153	6	3.9	0.24	0.1-0.6	9.4	0.002
Age groups						
6-12 m n=87	11	12.6	2	0.9-4.6	2.6	0.1
13-24 m n=103	9	8.7	1	0.4-2.4	0.004	0.95
25-36 m n=45	3	6.7	0.7	0.2-2.5	0.25	0.61
≥ 37 m n=44	1	2.3	0.21	0.02-1.6	2.66	0.102
Total	24	8.6	-	-	-	-

OR Odds ratio >1 (at risk)
 CI Confidence intervals
 χ^2 Chi-square ≥ 3.9 (significant)
 p Probability value ≤ 0.05 (significant)

Table 5: The associated odds ratio of susceptibility to polio virus infection with history of polio vaccine of malnourished children in Sana'a city Yemen.

Characters	Non-protective level N=24		OR	CI	χ^2	PV
	No	%				
First year vaccine						
Yes n=270	24	8.9		Undefined	0.87	0.34
No n=9	0	0		undefined	0.87	0.34
Booster dose						
Yes n=243	21	8.6	1.04	0.29-3.6	0.04	0.95
No n=36	3	8.3	0.96	0.27-3.4	0.4	0.95
National camping booster dose						
Yes n=183	10	5.5	0.49	0.21-1.13	6.8	0.01
No n=96	14	14.6	2.8	1.2-6.6	6.8	0.01

OR Odds ratio >1 (at risk)
 CI Confidence intervals
 χ^2 Chi-square ≥ 3.9 (significant)
 p Probability value ≤ 0.05 (significant)

Table 6: The associated odds ratio of susceptibility to polio virus infection with site of delivery, history of illness during first month of life, and type of feeding of malnourished children in Sana'a city Yemen.

Characters	Non-protective level N=24		OR	CI	χ^2	PV
	No	%				
Site of delivery						
Hospital n=165	15	9.1	1.2	0.49-2.7	0.12	0.72
House n=114	9	7.9	0.85	0.36-2.1	0.12	0.72
History of first month illness						
Yes n=158	15	9.5	1.4	0.6-3	0.36	0.54
No n=121	9	7.4	0.7	0.3-1.8	0.56	0.54
Type of feeding						
Mother feeding n=195	15	7.7	0.7	0.3-1.6	0.18	0.40
Bottle feeding n=75	6	8	0.89	0.34-2.3	0.05	0.82
Mixed n=9	3	33.3	5.9	1.4-25.4	7.2	0.007

OR Odds ratio >1 (at risk)
 CI Confidence intervals
 χ^2 Chi-square ≥ 3.9 (significant)
 p Probability value ≤ 0.05 (significant)

Reviewer's Copy