Traditional Chinese Medicine Endotherapy of Diabetic Wounds: A Scoping Review

Abstract

Background: In recent years, the incidence of diabetes has been increasing, and the diabetic foot is a complication of diabetes, which seriously affects the life and health of patients. The effect of Chinese medicine in the treatment of diabetes is more and more recognized.

Objectives:To search for the treatment of diabetic foot using Traditional Chinese medicine - Just to collect research literature on TCM treatment of diabetes in order to summarize the common syndrome types and medication rules in recent years, and to provide a new treatment.

Results: Common syndrome has syndrome of the dual deficiency of qi and yin, syndrome of dual deficiency of qi and yin, syndrome of meridian stasis-heat, syndrome of meridian stasis-heat and so on. Common endotherapy medicine is clear heat medicinal, tonify qi medicine, activate blood and resolve stasis and so on.

Conclusion: Traditional Chinese medicine has certain advantages and effects in the treatment of diabetes, but there is still no uniform standard

Keywords: Diabetic wounds, Traditional Chinese medicine, Endotherapy

Introduction

Diabetic foot secondary to the diabetic peripheral vascular disease and peripheral neuropathy caused by foot infection, ulcers or deep tissue destruction, is one of the more common serious complications in patients with diabetes. With the incidence rate of 6% - 20%, about 15% of diabetic foot patients will have foot ulcers or gangrene, which is an important cause of disability and death of patients with diabetes. [1]

With the aging of the population, the rising incidence of diabetes and the prolongation of the existing patients with type 2 diabetes, the incidence of diabetic lower extremity vascular disease is significantly increased. Diabetic foot treatment is difficult, with a high disability rate, and serious patients even need Amputation. According to the dialectical classification of TCM, the diabetic foot can be divided into main symptoms of blood stasis obstructing the vessel, lower energizer dampness-heat, blood stagnation due to the cold congealing. Traditional Chinese medicine has unique curative effects for different types of syndromes.[2,3]

In recent years, research on the treatment of diabetic foot by Chinese medicine has gradually increased, but there are few studies on common typesand common medications. In this paper, the scoping review method is used to screen, evaluate and summarize the research on Traditional Chinese Medicine internal treatment of diabetic foot, in order to further explore the common syndrome types and medication rules of Traditional Chinese Medicine treatment of diabetes, and provide new ideas for clinical.

Methodology

The methodology for this scoping review was based on the framework outlined by Arksey

and O'Malley (2005) and ensuing recommendations made by Levac *et al.* (2010). [4]This review is to collect research literature on TCM treatment of diabetes in order to summarize the common syndrome types and medication rules in January, 2015-November, 2019.

Aims:

The purpose of this study is to collect different methods of Traditional Chinese Medicine treatment of diabetic foot. This study focused on the analysis of common types and common medications used by Chinese medicine to treat diabetic foot. The study also collected the number of articles published in a specific time period (Figure 2). This article also conducts frequent statistics on the use of prescriptions and Traditional Chinese Medicine syndromes.

Search Strategy:

A systematic literature review was performed according to the PRISMA Statement.[5]CNKI(China National Knowledge Infrastructure) was this study's search engine, using the (allintitle:Diabetic foot AND Chinese medicine) The duration of the studies included was between 1st January 2015 till 1 st November 2019.

Data Extraction:

Using the CNKI search engine, the record was identified (n=58). Irrelevant articles (n=44) were removed. The records screened (n=33) because of some articles like Chinese medicine prevention of disease, external treatment of Chinese medicine, screeningarticles, abstract only, reviews articles, research progress articles, articles on method development, duration of study after November 2019 is out of bounds. Then led to the investigation of eleven (n=11)articles for the qualitative synthesis. The results have been shown in a Prisma flow diagram (Figure 1).

Result and Discussion

Diabetic peripheral neuropathy (DPN) is one of the common chronic complications of diabetes, and lower limbs are more serious than upper limbs, which are closely related to diabetic foot ulcers. The pathogenesis is complicated, and it is not clear at present. There is no specific treatment in western medicine. Western medicine has not only unsatisfactory efficacy, but also limited clinical application due to its high cost and side effects. In clinical studies, the dialectical treatment of this disease with traditional Chinese medicine (TCM) classification has achieved a good therapeutic effect and the side effects of traditional Chinese medicine treatment are small. Compared with western medicine, traditional Chinese medicine has the advantage of comprehensive regulation in the treatment of diabetic peripheral neuropathy.[6,7]

In this paper, the classification and frequency statistics of TCM syndromes and drug use rules of 11 included literatures were conducted using tables, and the analysis results are summarized as follows.

TCM Syndrome type

Among the 11 literatures, 10 mentioned the syndrome types of diabetic foot, with a total of 13 syndromes, with a cumulative frequency of 31. The top five with high frequency are: (1) syndrome of the dual deficiency of qi and yin, (2) syndrome of yin deficiency with congealing phlegm, (3) syndrome of dampness-heat obstruct, (4) syndrome of qi deficiency with blood stasis, (5) syndrome of meridian stasis-heat. See Table 1.[1-3,6-13] Table 1 shows that the syndromes of diabetic foot are heterogeneous and variable, and they are always sign of deficiency. This deficiency mainly includes deficiency of qi and blood and a deficiency of qi and Yin. Standards include blood stasis, heat, phlegm and so on. At

the beginning of the main virtual; In the middle stage, due to the invasion of external evil and internal evil, the righteous spirit still exists, and the righteous and evil are competing with each other. Due to the long course of the disease, consumption injury qi will convert to virtual at last.

Drug of TCM

Applied Chinese medicines

Among the 11 selected literatures, a total of 84 TCM varieties were used for TCM syndrome differentiation treatment, with a cumulative frequency of 140. The results showed that the top 6 TCM medicines were respectively: Salvia miltiorrhiza (5/140), Astragalus(5/140), Angelica(5/140), prepared rehmannia root(4/140), Safflower(4/140), Ligusticum striatum(4/140). Their application rate is 3.57%, 3.57%, 3.57% and 2.86% respectively. See Table 2..[1-3,6-13]

Traditional Chinese medicine efficacy

According to the statistics of The Times of using traditional Chinese medicine with different efficacy in the treatment of diabetes in 11 literatures, the results showed that The Times of using traditional Chinese medicine ranked the top five respectively: medicinal of clear heat(19/84), medicinal of tonify qi(16/84), medicinal of activate blood and resolve stasis(14/84), medicinal of tonify yin(10/84), medicinal of regulate qi(8/84). Their application rate is 22.62%, 19.05%, 16.67%, 11.90%, 9.52% respectively. See Table 3.[1-3,6-13]

Conclusion

The treatment of clinical diabetes mainly includes Western medicine treatment and Traditional Chinese medicine treatment. Western medicine takes effect quickly but need to take hypoglycemic drugs for a long time. The illness will come back easily and it has relatively large side effects, such as dizziness, cough and so on. According to TCM theory, diabetic foot ulcer is caused by deficiency of qi and Yin, and the main syndrome is the deficiency of qi and blood stasis, which is marked by excess of dampness and heat. Therefore, the treatment and nursing of diabetic foot ulcer should adopt the treatment methods of activating blood circulation and removing blood stasis, clearing heat and detoxification and the nursing methods of internal and external conditioning of Traditional Chinese medicine, pay attention to the local nursing of affected foot, so as to make the skin of foot turn red, reduce pain and promote ulcer healing. However, we still need to develop a more scientific classification method for TCM syndromes of diabetic foot.

In conclusion, active TCM syndrome differentiation treatment for diabetic foot ulcer patients can effectively control the clinical symptoms of patients, improve the treatment effect and improve the prognosis of patients, which are worthy of clinical application and promotion.

Reference:

- 1. Zhou G. Y. & Zhang W.K. (2015). Preliminary comparison of curative effect of traditional Chinese medicine and western medicine in treating diabetic foot. Journal of Microcirculation, 25 (01), 34-36.
- 2. Cheng Y. Y. (2015). Observation on the curative effect of integrated traditional Chinese medicine and traditional Chinese medicine on early diabetic foot (stagnation and obstruction). Chinese Journal of Integrated Traditional and Western Medicine on Cardiovascular Disease, 3 (24), 100-101."
- 3. Zhang A. M. (2017). Treatment of 160 Cases of Diabetic Lower Extremity Vascular

- Diseases with Internal and External Therapy of Traditional Chinese Medicine. Modern Distance Education of Chinese Medicine, 15 (03), 58-60 + 67."
- 4. Pham, M. T., Rajić, A., Greig, J. D., Sargeant, J. M., Papadopoulos, A., & McEwen, S. A. (2014). A scoping review of scoping reviews: advancing the approach and enhancing the consistency. Research synthesis methods, 5(4), 371-385.
- 5. Jin, K. Y., & Aslam, M. S. (2019). The occurrence of pharmaceutical waste in different parts of the world: A scoping review. PeerJ Preprints, 7, e27951v1.
- 6. Deng S. G. & Wu M. Q. (2017). Liao Fang. Clinical observation of Huoxue Tongmai Decoction in treating diabetic foot. Clinical Research of Traditional Chinese Medicine, 9 (07), 56-57 + 59.
- 7. Gong G. Y. (2017). Clinical Observation on 36 Cases of Comprehensive Treatment of Diabetic Foot in Community. China Medical Journal of Metallurgical Industry, 34 (01), 85.
- 8. Liu A. X. (2016). Clinical effect of comprehensive treatment of traditional Chinese medicine and western medicine on elderly patients with type 2 diabetic peripheral neuropathy. Clinical Research of Traditional Chinese Medicine, 8 (35), 68-69.
- 9. Liu C., Wang J. & Hou J. (2019). Discussion on clinical effects of TCM syndrome differentiation on diabetic foot. Biped and Health, 28 (10), 49-50.
- 10. Peng G. H.& Liu F. F. (2015). Observation on Traditional Chinese Medicine Effect of Peripheral Neuropathy of Diabetes with Cold Coagulation and Stasis Type. Shenzhen Journal of Integrated Traditional Chinese and Western Medicine ,25 (11), 35-37.
- 11. Wang Y. L., Liao M. J., Li Y. & Tong G. (2019). Therapeutic effect of external therapy of traditional Chinese medicine combined with Yanghe Decoction on diabetic foot wounds. Electronic Journal of Foot and Ankle Surgery, 6 (03), 25-26 + 29.
- 12. Xiong W. (2019). Efficacy of Integrated Traditional Chinese and Western Medicine in Treating Non-Ischemic Diabetic Foot. Journal of Clinical Medical Literature, 6 (06), 11.
- 13. Yang H. W. (2018). Exploring the clinical treatment effect of diabetic foot patients with comprehensive treatment of traditional Chinese and western medicine. World Medical Digest, 18 (65), 170 + 172

Table 1: Frequency of TCM Syndrome type

| | Equency of 1 CM Syndrome type | D | (a) |
|----------------|---|----------|---|
| The | TCM Syndrome type | Frequenc | Rate (%) |
| sequence | | y | |
| 1 | syndrome of dual deficiency of qi and yin | 7 | 22.58 Records |
| 2 | syndrome of yin deficiency with | 5 | 16.13 excludedChinese |
| | congealing phlegm | | medicine prevention of |
| 3 | syndrome of dampness-heat obstructing | 3 | 9.68 disease, external |
| 4 | syndrome of qi deficiency with blood | 3 | 9.68 treatment of Chinese |
| | stasis | | medicine, screening |
| 5 | syndrome of meridian stasis-heat | 3 | 9.68 articles ,abstract |
| Identi | | | only ,reviews articles, research progress articles, articles on |
| Identification | Records identified throug n =58 | gh CNKI | method development, Duration of study after November 2019 n =11 |




Figure1Prisma Flow Diagram




Figure2 Number of Articles in the Study That WerePublished Between January 2015-November 2019

Table 2: Frequency of Applied Chinese medicine

| The sequence | Name of drug | Frequency Rate (%) |
|--------------|-------------------------|--------------------|
| 1 | Salvia miltiorrhiza | 5 3.57 |
| 2 | Astragalus | 5 3.57 |
| 3 | Angelica | 5 3.57 |
| 4 | prepared rehmannia root | 4 2.86 |
| 5 | Safflower | 2.86 |
| 6 | Ligusticum striatum | 4 2.86 |

Table 3: Frequency of Traditional Chinese medicine efficacy

| The sequence | Name of drug | Frequency | Rate (%) |
|--------------|---|-----------|----------|
| 1 | medicine of clear heat | 19 | 22.62 |
| 2 | medicine of tonify qi | 16 | 19.05 |
| 3 | Medicine of activate blood and resolve stasis | 14 | 16.67 |
| 4 | medicine of tonify yin | 10 | 11.90 |
| 5 | medicine of regulate qi | 8 | 9.52 |