



## RESEARCH ARTICLE

## EVALUATION OF THE FREE RADICAL SCAVENGING OF TECTOQUINONE COMPOUND ISOLATED FROM *SYZYGIUM OBLANCEOLATUM* (C.B.ROB.)

MERR

Ahmad Najib , Virsa Handayani , Muhammad Rizki Jaelani

Laboratory of Pharmacognosy-Phytochemistry, Phytochemistry Division, Faculty of Pharmacy, Universitas Muslim Indonesia, Indonesia.

## Article Info:



## Article History:

Received: 1 August 2023  
 Reviewed: 10 September 2023  
 Accepted: 27 October 2023  
 Published: 15 November 2023

## Cite this article:

Najib A, Handayani V, Jaelani MR. Evaluation of the free radical scavenging of tectoquinone compound isolated from *Syzygium oblanceolatum* (C.B.Rob.) Merr. Universal Journal of Pharmaceutical Research 2023; 8(5):10-14.

<https://doi.org/10.22270/ujpr.v8i5.1003>

## \*Address for Correspondence:

Ahmad Najib, Laboratory of Pharmacognosy-Phytochemistry, Phytochemistry Division, Faculty of Pharmacy, Universitas Muslim Indonesia, Indonesia, Tel: +62 815-2404-5514  
 E-mail: [ahmad.najib@umi.ac.id](mailto:ahmad.najib@umi.ac.id)

## Abstract

**Aim and objective:** In this study, the free radical scavenging of tectoquinone compound isolated from leaf of *Syzygium oblanceolatum* (C.B. Rob.) Merr was investigated by assessing their capacity to scavenge free radicals using the DPPH (2, 2-diphenyl-1-picrylhydrazyl) assay.

**Methods:** The quantitative assessment of antioxidant activity was conducted using a UV-Vis spectrophotometric approach, with measurements taken at a wavelength of 516 nm.

**Results:** The IC<sub>50</sub> value, representing the concentration required to inhibit 50% of the DPPH radicals, was determined to be 66.362 µg/mL, indicating a moderate free radical-scavenging activity.

**Conclusion:** These findings suggest that tectoquinone compounds possess a discernible ability to against free radical, though further research may be necessary to optimize their potential applications in health and medicine.

**Keywords:** DPPH assay, free radical, IC<sub>50</sub> value, *Syzygium oblanceolatum*, Tectoquinone.

## INTRODUCTION

Oxidative stress is a fundamental contributor to various health disorders and is primarily associated with the detrimental effects of reactive oxygen species (ROS) on biological systems<sup>1</sup>. Consequently, the exploration of natural compounds with antioxidant properties has become a focal point in scientific research, as these compounds have the potential to mitigate the adverse effects of free radical compounds<sup>2</sup>.

Tectoquinone compound, isolated from the plant *Syzygium oblanceolatum* (C.B. Rob.) Merr, represents a group of compounds that have attracted attention for their potential as free radical scavenging. *S. oblanceolatum* represents the first documented report of its existence in Sulawesi, previously known to occur only in Eastern Philippines and Kalimantan<sup>3</sup>. The presence of this plant in Sulawesi, particularly in South Sulawesi, is intriguing and warrants further investigation. *S. oblanceolatum* from a family of Myrtaceae is a well-known family of vascular dicot plants<sup>4</sup>. As the eighth-largest dicot plant family, it

comprises approximately 5,650 species grouped into 130-150 generad<sup>5</sup> and is widely distributed across Africa, extending into South Asia and tropical Southeast Asian countries<sup>6</sup>. Due to their phytochemical content and health-promoting properties, several *Syzygium* species have garnered significant interest<sup>7,8</sup>. On the other hand, many other *Syzygium* species remain relatively unexplored<sup>9</sup>, and their chemical and biological activities continue to be of interest including the class of quinones compound<sup>10</sup>.

Quinones, a class of organic compounds, exhibit diverse medical properties<sup>11</sup>. They are recognized for their antioxidant capabilities<sup>12</sup>, with compounds like Coenzyme Q10 and Vitamin K helping combat oxidative stress<sup>13</sup>. In cancer treatment, specific quinones, such as anthracyclines, interfere with cancer cell growth<sup>14</sup>. Quinones also play a role in skincare, blood clotting<sup>15</sup>, and neurological disorders<sup>16</sup>. Some quinones demonstrate antibacterial<sup>17</sup>, antifungal<sup>18</sup>, and anti-inflammatory effects<sup>19</sup>. CoQ10, in particular, is studied for mitochondrial function and cardiovascular health<sup>20</sup>. While promising, the use of quinone-based

treatments should always involve consultation with healthcare professionals due to potential side effects and interactions with other medications.

Tectoquinone (as a member of the class of quinones compound)<sup>21</sup> is a naturally occurring compound that has attracted attention due to its potential antioxidant properties. It is often found in certain plant species, and researchers have been investigating its ability to scavenge free radicals, which are harmful molecules associated with oxidative stress and various health issues. Tectoquinone is a type of organic compound. Specifically, it belongs to the class of compounds known as quinones. Quinones are a class of organic compounds characterized by a six-membered aromatic ring containing two carbonyl (C=O) functional groups<sup>22</sup>. Tectoquinone, like other quinones, is known for its chemical structure that includes this characteristic aromatic ring with carbonyl groups, which contributes to its reactivity and potential biological activities, including its antioxidant properties<sup>23</sup>.

In this study, we investigate the capacity of tectoquinone compounds to scavenge free radicals, a critical aspect of their antioxidant activity. The widely recognized DPPH (2,2-diphenyl-1-picrylhydrazyl) assay was employed for this purpose, offering a reliable means to assess the radical-scavenging abilities of compounds.

To provide a quantitative assessment of the free radical scavenging activity, a UV-Vis spectrophotometric approach was utilized, enabling precise measurements at a specific wavelength. This method allowed us to obtain valuable data on the antioxidant potential of the tectoquinone compound.

## MATERIALS AND METHODS

### Materials

The samples used in this research are tectoquinone compound isolated from leaf of *Syzygium oblongatum* (C.B.Rob.) Merr. was collected in Maros district, South Sulawesi-Indonesia. The collected plant was identified by a taxonomist, Dr. Wu-Kuang Soh (Trinity College Dublin, the Republic of Ireland), DPPH (Sigma Aldrich), Quercetin (Sigma Aldrich).

### Methods

The study was carried out through experimental procedures conducted in a laboratory setting, employing the free radical scavenging method. Tectoquinone previously identified by NMR Spectroscopy and compared with literature.

### Sample extraction and isolation

The leaves of *S. oblongatum* (C.B.Rob.) Merr. ground into a powder (0.7 kg) were extracted three times with 20 L each of MeOH by maceration at room temperature. The filtrates were combined and evaporated under reduced pressure to yield 60 g of a dark gummy extract. The extract (10.1 g) was suspended in methanol: water (9:1) (0.2 L) and extracted with *n*-hexane (7 x 0.25 L) to obtain the *n*-hexane extract 2.7 g and methanol: water 7.3 g. The *n*-hexane extract was applied to a silica gel column and eluted with an *n*-hexane, *n*-hexane: ethyl acetate (8:1;

6:1; 4:1; 1:1) and methanol respectively to give seven major fractions (H1–H7). Fraction H2 (340.5 mg) was applied to a silica gel column and eluted with *n*-hexane: acetone (15:1) and methanol to yield nine fractions (H2A–H2I). Fraction H2H1 was separated successively by preparative reversed-phase HPLC using the eluent methanol: water (3:1) with flow rate 8 mL/min to afford three fractions (H2H1A–H2H1C). Fraction H2H1B was further purified by reversed-phase preparative thin layer chromatography PTLC eluted with methanol: water (5:1) to yield compound tectoquinone (2.0 mg).

### Preparation of DPPH stock solution

To prepare a DPPH solution with a concentration of 30 ppm, 1.5 mg of DPPH powder was dissolved in 50 mL of high-quality analytical methanol within a volumetric flask. Following this, the measurement of the DPPH's maximum wavelength was conducted within the range of 450 nm to 550 nm.

### Preparation Quercetin standard solution

To prepare a reference quercetin solution, a concentration of 1,000 ppm was initially established. Subsequently, various concentrations, including 0.2 ppm, 0.4 ppm, 0.6 ppm, 0.8 ppm, and 1 ppm, were derived from this primary solution, with each variant comprising a 5 mL volume. A 2 mL aliquot of each quercetin solution series was then measured and transferred to separate vials. Next, 2 mL of DPPH solution was introduced into each vial. Following thorough mixing, the vials were left to incubate in a dark environment for 30 minutes. After this incubation period, the absorption at the maximum wavelength was determined using a UV-Vis spectrophotometer, namely 516 nm.

### Preparation of Tectoquinone

Weighing 0.7 mg of the tectoquinone compound, we then dissolved it in 2.5 mL of high-quality analytical methanol, resulting in a stock solution with a concentration of 280 ppm. Following that, a range of concentrations was created by diluting 4 mL of this solution to produce concentrations of 10 ppm, 20 ppm, 30 ppm, 40 ppm, and 50 ppm.

### Examination of free radical scavenging

To assess the radical scavenging of tectoquinone compound, 2 mL of test solutions with concentrations ranging from 10 ppm to 50 ppm were mixed with an equal volume of DPPH solution in vials and left to incubate in darkness for 30 minutes. Following this incubation period, their absorbance was measured at the maximum wavelength, namely 516 nm.

The percentage of DPPH radical inhibition was determined using the formula:

$$\% \text{ inhibition} = \frac{A_1 - A_2}{A_1} \times 100$$

Where A1 represents the absorbance of the control and A2 denotes the absorbance of the sample.

To find the IC<sub>50</sub> value, a linear curve was established by plotting the test solution concentrations (x-axis) against the corresponding % inhibition (y-axis) using the equation  $y = a + bx$ . The IC<sub>50</sub> value was then calculated as  $IC_{50} = (50 - a)/b$ .

## RESULTS AND DISCUSSION

The  $^1\text{H}$  NMR spectrum showed five protons coupling with one another at  $\delta_{\text{H}}$  7.47-7.36 (m, 5H) indicating the absence of an aromatic ring. The signal at  $\delta_{\text{H}}$  5.40 (dd,  $J=12.7, 3.3$  Hz, 1H) showed correlation with a geminal protons at  $\delta_{\text{H}}$  3.04 (dd,  $J = 17.0, 12.9$  Hz, 1H), and  $\delta_{\text{H}}$  2.84 (dd,  $J=17.4, 2.9$  Hz, 1H). In addition, broad singlet for two methyl groups  $\delta_{\text{H}}$  2.07 and 2.06 (d,  $J=3.1$  Hz, 6H), one of the protons in the downfield area at  $\delta_{\text{H}}$  12.26 indicated an absence of hydroxyl proton. The  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ) showed 15 signals: three signals due to the oxygen substituted aromatic carbon at  $\delta_{\text{C}}$  160.86, 159.39, 157.72, seven signals due to carbon or hydrogen substituted aromatic at  $\delta_{\text{C}}$  139.02, 128.87, 128.66, 125.97, 103.05, 102.94, 102.06 and a signal each oxygen substituted carbon at  $\delta_{\text{C}}$  78.77, methylene at  $\delta_{\text{C}}$  43.56, carbonyl carbon at  $\delta_{\text{C}}$  196.42 and two methyl carbon at  $\delta_{\text{C}}$  7.7 and 6.9. Spectrum data were compared with previous literature isolated compound was found as tectoquinone<sup>24</sup>.


Tectoquinone: pale yellow solid;  $[\alpha]_{\text{D}}^{23}=-72.81$  ( $c=0.070, \text{CHCl}_3$ );  $\text{DART}^+m/z$  285.042  $[\text{M}+1]^+$  (Cal  $\text{C}_{17}\text{H}_{16}\text{O}_4$  285.105  $[\text{M}+1]^+$ );  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  12.26 (s, 1H), 7.47-7.36 (m, 5H), 5.40 (dd,  $J=12.7, 3.3$  Hz, 1H), 3.04 (dd,  $J = 17.0, 12.9$  Hz, 1H), 2.84 (dd,  $J = 17.4, 2.9$  Hz, 1H), 2.07 (d,  $J = 3.1$  Hz, 6H) and  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  196.42 (C-4), 160.86 (C-7), 159.39 (C-5), 157.72 (C-9), 139.02 (C-1'), 128.87 (C-3'; C-5'), 128.66 (C-4'), 125.97 (C-2'; C-6'), 103.05 (C-10), 102.94 (C-8), 102.06 (C-6), 78.77 (C-2), 43.56 (C-3), 7.7, 6.9

In the quantitative assessment of antioxidant potential, the DPPH scavenging method relies on the  $\text{IC}_{50}$  value. This value indicates the concentration of the test sample required to achieve a 50% inhibition of oxidative processes (effectively reducing or inhibiting oxidation by 50%). The outcomes of absorbance measurements, percentage inhibition, and  $\text{IC}_{50}$  values for both tectoquinone compound and the reference quercetin are provided in the Table 1.

**Table 1: Results of absorbance measurements, percentage inhibition, and  $\text{IC}_{50}$  values.**

Sample	Concentration (ppm)	Blank Absorbance	Sample Absorbance	Inhibition (%)	$\text{IC}_{50}$ ( $\mu\text{g}/\text{mL}$ )
Tectoquinone	10	0.596	0.346	41.750	66.362
	20	0.596	0.335	43.602	
	30	0.596	0.321	45.959	
	40	0.596	0.319	46.296	
	50	0.596	0.313	47.306	
Quercetin	0.2	0.596	0.310	47.986	0.237
	0.4	0.596	0.264	55.704	
	0.6	0.596	0.252	57.718	
	0.8	0.596	0.245	58.892	
	1.0	0.596	0.204	65.771	

Based on the data above the relationship between % inhibition and the concentration of tectoquinone compounds is depicted in Figure 1. Linear regression equations were established on the graphs, where concentration is plotted on the x-axis, and % inhibition is on the y-axis. Consequently, the  $\text{IC}_{50}$  value for tectoquinone compounds can be determined from the regression equation.



**Figure 1: Relationship between tectoquinone concentrations and % inhibition.**

The linear regression obtained from the quercetin reference is  $y=3.8758x+45.587$  with an  $R^2$  value of 0.9175, and for tectoquinone compounds, it is  $y = 0.138x + 40.842$  with an  $R^2$  value of 0.9329. These

equations can then be rearranged into the form  $y=bx + a$ , where  $y$  represents 50% inhibition, and  $x$  is the  $\text{IC}_{50}$  value.

A compound is considered a very strong antioxidant if the  $\text{IC}_{50}$  value is  $<10 \mu\text{g}/\text{mL}$ , strong if it falls between  $10-50 \mu\text{g}/\text{mL}$ , moderate if it ranges between  $50-100 \mu\text{g}/\text{mL}$ , weak if it falls between  $100-250 \mu\text{g}/\text{mL}$ , and inactive if the  $\text{IC}_{50}$  value is above  $250 \mu\text{g}/\text{mL}$ <sup>25</sup>. From the results obtained in Table 1, the  $\text{IC}_{50}$  value for the quercetin reference is  $0.237 \mu\text{g}/\text{mL}$ , classifying it as a very strong antioxidant because it falls within the  $\text{IC}_{50}$  range of  $<10 \mu\text{g}/\text{mL}$ . Meanwhile, the  $\text{IC}_{50}$  value for tectoquinone compounds is  $66.362 \mu\text{g}/\text{mL}$ , categorizing it as a moderate antioxidant within the  $50-100 \mu\text{g}/\text{mL}$  range. As explained previously mention that tectoquinone belongs to the quinone class compound. Quinone compounds inhibit free radicals through their unique redox properties, enabling them to participate in electron transfer reactions.

When quinones encounter free radicals, they donate electrons to neutralize these highly reactive species, rendering them less harmful. This process transforms quinones into semiquinone radicals, which are relatively stable and less reactive. In some cases, semiquinone radicals can further react to regenerate the original quinone molecule, allowing quinones to continue their antioxidant action through multiple redox cycles.

This mechanism makes quinones effective anti-oxidants, protecting cells and biomolecules from the damaging effects of oxidative stress caused by free radicals.

#### Limitations of the study

The limitation of this study is that it primarily discusses the general properties and mechanisms of quinone compounds in inhibiting free radicals, with a specific focus on tectoquinone. However, it lacks specific experimental data or results related to tectoquinone's antioxidant activity or its comparison with other known antioxidants. While the theoretical background on quinones is valuable, the study does not provide empirical evidence of tectoquinone's antioxidant efficacy. To draw meaningful conclusions about tectoquinone's potential as an antioxidant, further experimental research, including *in vitro* or *in vivo* assays, is required to assess its actual free radical-scavenging capacity and compare it with established antioxidants. Additionally, the study does not address potential side effects, bioavailability, or practical applications of tectoquinone, which are essential considerations for its use in health and medicine.

#### CONCLUSION

The results indicate that tectoquinone compounds exhibit a noticeable capability against free radicals with moderate potential activity; however, additional research might be needed to enhance their potential uses in the field of health and medicine.

#### ACKNOWLEDGEMENTS

The author would like to extend their thanks to the Research and Development Institute of Resources (LP2S) at the Muslim University of Indonesia (UMI) for financially supporting this study. Additionally, gratitude is also expressed to Associate Professor Kyoko Nakagawa-Goto from the Laboratory of Natural Products and Medicinal Chemistry at Kanazawa University, Japan, for granting access to NMR facilities for the identification of tectoquinone compounds.

#### AUTHOR'S CONTRIBUTION

**Najib A:** writing original draft, methodology, investigation. **Handayani V:** writing, review and editing. **Jaelani MR:** formal analysis, methodology. Every author gave their approval to the manuscript's final draft.

#### DATA AVAILABILITY

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

#### CONFLICT OF INTEREST

No conflict of interest is associated with this work.

#### REFERENCES

- Anik MI, Mahmud N, Masud AAI, *et al.* Role of reactive oxygen species in aging and age-related diseases: A review. *ACS Appl Bio Mater* 2022;5(9):4028–54. <https://doi.org/10.1021/acsabm.2c00411>
- Jia W, Wu X, Kang X. Integrated the embedding delivery system and targeted oxygen scavenger enhances free radical scavenging capacity. *Food Chem X* 2023;100558. <https://doi.org/https://doi.org/10.1016/j.fochx.2022.100558>
- Ashton P. Notes on northern Borneo Myrtaceae. *Sandakania*. 2016; 22:55–75.
- Armijos C, Ramirez J, Vidari G. Poorly investigated ecuadorian medicinal plants. *Plants* 2022; 11(12):1590. <https://doi.org/10.3390/plants11121590>
- Rocchetti G, Lucini L, Ahmed SR, Saber FR. *In vitro* cytotoxic activity of six *Syzygium leaf* extracts as related to their phenolic profiles: An untargeted UHPLC-QTOF-MS approach. *Food Res Int* 2019;126(August):108715. <https://doi.org/10.1016/j.foodres.2019.108715>
- Pham NKT, Nguyen TD, Doan TDC, *et al.* Stercufotoin A, new oleanane-type triterpenoid from the leaves of *Sterculia foetida* L. *Nat Prod Res* 2021;35(7):1226–31. <https://doi.org/10.1080/14786419.2019.1644508>
- Kumar M, Zhang B, Nishad J, *et al.* Jamun (*Syzygium cumini* (L.) Skeels) Seed: A review on nutritional profile, functional food properties, health-promoting applications, and safety aspects. *Processes*. 2022;10(11):2169. <https://doi.org/https://doi.org/10.3390/pr10112169>
- Amir Rawa MS, Mazlan MKN, Ahmad R, Nogawa T, Wahab HA. Roles of syzygium in anti-cholinesterase, anti-diabetic, anti-inflammatory, and antioxidant. *Alzheimer's Persp Plants* 2022;11(11):1476. <https://doi.org/10.3390/plants11111476>
- Pesiu E, Lee GE, Salam MR, *et al.* Species composition, diversity, and biomass estimation in coastal and marine protected areas of Terengganu, Peninsular Malaysia. *Agronomy*. 2022; 12(10):2380. <https://doi.org/10.3390/agronomy12102380>
- de Sousa DP, Damasceno ROS, Amorati R, *et al.* Essential oils: Chemistry and pharmacological activities. *biomolecules*. 2023;13(7):1144. <https://doi.org/10.3390/biom13071144>
- Dahlem Junior MA, Nguema Edzang RW, Catto AL, Raimundo JM. Quinones as an efficient molecular scaffold in the antibacterial, antifungal or antitumoral arsenal. *Int J Mol Sci* 2022;23(22):14108. <https://doi.org/10.3390/ijms232214108>
- Prokopiev I, Sleptsov I, Serebryakov E, Sharoyko V. Antioxidant and cytotoxic activities of quinones from *Cetraria laevigata*. *Nat Prod Res* 2023:1–5. <https://doi.org/10.1080/14786419.2023.2187795>
- Cores A, Carmona-Zafra N, Clerigué J, Villacampa M, Menéndez JC. Quinones as Neuroprotective Agents. *Antioxidants* 2023; 12(7):1464. <https://doi.org/10.3390/antiox12071464>
- Kiran P, Ghosh A, Pawar V, Maske P, Khan A, Srivastava R. Hypoxia-targeting drugs as new cancer chemotherapy agents: Molecular Insights. *Hypoxia in cancer: Significance and impact on cancer therapy*. Springer 2023; 351–68. [https://doi.org/10.1007/978-981-99-0313-9\\_16](https://doi.org/10.1007/978-981-99-0313-9_16)
- Mishima E, Wahida A, Seibt T, Conrad M. Diverse biological functions of vitamin K: from coagulation to ferroptosis. *Nat Metab*; 2023:1–9. <https://doi.org/10.1038/s42255-023-00821-y>
- Peng Y, Xu D, Mao S, Zhou X. Neurotoxicity and apoptosis induced by pyrroloquinoline quinone and its ester derivative on primary cortical neurons. *Neurotoxicol* 2020;78:47–56. <https://doi.org/10.1016/j.neuro.2020.02.005>
- Sahoo PMS, Behera S, Behura R, *et al.* A brief review: Antibacterial activity of quinone derivatives. *Biointerface Res Appl Chem* 2022; 12:3247–58. <https://doi.org/10.33263/BRIAC123.32473258>

18. Pacios-Michelena S, González CNA, Herrera RR, *et al.* Biomass from phytopathogens and culture conditions improve *Penicillium chrysogenum* antimicrobial activity and antifungal compounds production. *Environ Qual Manag* 2023;33(1):349–58. <https://doi.org/10.1002/tqem.22065>
19. Mohamad Ishak NS, Ikemoto K. Pyrroloquinoline-quinone to reduce fat accumulation and ameliorate obesity progression. *Front Mol Biosci* 2023; 10:1200025. <https://doi.org/10.3389/fmolb.2023.1200025>
20. Pu TT, Wu W, Liang PD, Du JC, Han SL, Deng XL, *et al.* Evaluation of coenzyme Q10 (CoQ10) deficiency and therapy in mouse models of cardiomyopathy. *J Cardiovasc Pharmacol* 2022:10–1097. <https://doi.org/10.3390/antiox11091698>
21. Martin JA, López R. Biological deterioration and natural durability of wood in Europe. *Forests* 2023;14(2):283. <https://doi.org/10.3390/f14020283>
22. Zhang S, Zhu YL, Ren S, *et al.* Covalent Organic Framework with multiple redox active sites for high-performance aqueous calcium ion batteries. *J Am Chem Soc* 2023; 145(31):17309–20. <https://doi.org/10.1021/jacs.3c04657>
23. Cárcamo-Ibarra E, Manuel Martinez-Pacheco M, Munro-Rojas A, *et al.* Antifungal activity of crude extracts of *Tectona grandis* Lf against wood decay fungi. *Phyt* 2022; 91(8). <https://doi.org/10.32604/phyton.2022.020578>
24. Peralta MA, Santi MD, Agnese AM, Cabrera JL, Ortega MG. Flavanoids from *Dalea elegans*: Chemical reassignment and determination of kinetics parameters related to their anti-tyrosinase activity. *Phytochem Lett* 2014; 10:260–7. <https://doi.org/10.1016/j.phytol.2014.10.012>
25. Dahlia AA, Daim M, Najib A. Comparative study of antioxidant activity of turmeric ethanol extract (*Curcuma longa* L.) and ethanol extract of curcuma (*Curcuma xanthorrhiza* Roxb.) using the 1,1-diphenyl-2-picrylhydrazil (DPPH) radical scavenging method. *AIP Conference Proceedings* 2023. <https://doi.org/10.1063/5.0123987>