



RESEARCH ARTICLE

COMPARATIVE ANTI-TRICHOMONAS VAGINALIS ACTIVITY EVALUATION OF SAMBUCUS NIGRA L. FLOWERS AND FRUITS EXTRACTS

Ecem Erdemir¹ , Husniye Kayalar^{2*} , İbrahim Cavus³ , Ahmet Ozbilgin³

¹Bozyaka Education Research Hospital, Izmir, Turkey.

²Ege University Faculty of Pharmacy Department of Pharmacognosy, Bornova, Izmir, Turkey.

³Manisa Celal Bayar University, Faculty of Medicine, Department of Medical Parasitology, Manisa, Turkey.

Article Info:



Article History:

Received: 23 March 2024

Reviewed: 6 May 2024

Accepted: 27 June 2024

Published: 15 July 2024

Cite this article:

Erdemir E, Kayalar H, Cavus İ, Ozbilgin A. Comparative anti-*Trichomonas vaginalis* activity evaluation of *Sambucus nigra* L. flowers and fruits extracts. Universal Journal of Pharmaceutical Research 2024; 9(3): 46-50. <http://doi.org/10.22270/ujpr.v9i3.1121>

*Address for Correspondence:

Dr. Husniye Kayalar, Ege University Faculty of Pharmacy Department of Pharmacognosy, Bornova, Izmir, Turkey. Tel: +905543986021; E-mail: husniye.kayalar@ege.edu.tr

Abstract

Background: *Sambucus nigra* or also known as Black elderberry or European elderberry is been used for the treatment and prevention of various diseases. The plant possesses health beneficial effects such as antimicrobial, antioxidant, antiulcerogenic, antidepressant, antidiabetic, antiviral and antiinflammatory. However, there is limited report on the antitrichomoniasis activity of *S. nigra*.

Objectives: The study aimed to comparatively analyze the *in vitro* trichomonocidal effects of various extracts prepared from both fruits and flowers of *S. nigra*. Half maximal inhibitory concentration (IC₅₀) and minimum parasiticide (MPC) values of the extracts against metronidazole-resistant *Trichomonas vaginalis* 50143 strain were determined by the liquid microdilution method using 96-well microplates.

Results: After 48 hrs incubation with parasites, the highest activities were observed for 70% hydroethanolic extract, chloroform extract and methanol extract of fruits with IC₅₀ values of 513, 531 and 566 µg/ml, respectively. Fruit extracts exhibited higher activity than flower extracts, and no activity was observed in methanol extract of flowers.

Conclusion: This study is the first comparative analysis of the anti-*Trichomonas vaginalis* activity of elderberry extracts. This study will serve as a reference source in isolating the components of elderberry fruits with antiparasitic activity.

Keywords: Metronidazole, plant extracts, *Sambucus nigra*, *Trichomonas vaginalis*, trophozoites.

INTRODUCTION

Sambucus nigra L., distributed in Europe, North Africa, West and Central Asia and North America, is one of the two species growing wildly in Turkey. This perennial shrub or 8-10 m tall tree with purplish fruit and creamy white flowers belongs to Adoxaceae family¹⁻³. *S. nigra* is also known as Black elderberry or European elderberry and believed to have been used since ancient times for the treatment and prevention of various diseases^{4,5}. In traditional Chinese medicine *S. ebulus* or known as dwarf elder is widely used in bone and joint disorders⁶. In Europe and African countries, *Sambucus* species are traditionally used in the treatment of respiratory diseases such as asthma and bronchitis, throats, sinusitis, herpes, neuralgia, epilepsy and for dental problems for bronchitis and stomach ache^{7,8}. In Turkey, in addition to external use for the treatment of rheumatism and wounds, the decoction or infusion prepared from flowers and fruits of *S. nigra* are used as diuretic, diaphoretic, immune system

booster and protective against cancer. The fruits are also used in the carpet and textile industry due to its colorful structure⁹⁻¹².

Numerous pharmacological activities have been conducted on the fruits of *S. nigra* and proven to possess health beneficials such as antimicrobial, antioxidant, antiulcerogenic, anti-depressant, antidiabetic, antiviral and anti-inflammatory^{8,13}. Due to various health benefits, the elderberry fruits are also included in the composition of jam, pie, ice cream, yoghurt, wine, tea and fruit juices⁶.

Trichomoniasis is one of the most common sexually transmitted diseases caused by *T. vaginalis* whose annual average prevalence is around 156 million in the world¹⁴. *Trichomonas* manifests itself with a number of symptoms, and different symptoms occur in men and women. This disease is also reported to be the causative agent of cervical cancer, pelvic inflammation and infertility¹⁵. Metronidazole is the most effective drug used in the treatment of trichomoniasis. Unfortunately, besides drug resistance, side effects

such as gastrointestinal disorders, taste problems, nausea, vomiting, vertigo, neutropenia, skin reactions, cloudy urine and fatigue are reported for metronidazole¹⁶⁻¹⁹. Due to serious side effects of 5-nitroimidazole derivatives, a more accessible source of drugs with fewer side effects are still being investigated from natural sources²⁰.

In a study conducted in Iran, plants registered in international databases were searched for their *in vitro* anti-*Trichomonas* activity. Garlic, lavender, eucalyptus, geranium, onion, yarrow, wormwood which are traditionally used for antiparasitic activities were found to be effective against *T. vaginalis*²¹. In other investigations conducted to find alternative sources for the treatment of trichomoniasis, rhubarb, turmeric and fennel were found to inhibit the growth and reproduction of parasites^{22,23}. In a recent study, Niknam *et al.*, reported the anti-*T. vaginalis* effect of methanolic extract prepared from *S. nigra* fruits from northern Iran²⁴.

To the best of our knowledge this is the first comparative research in scientific literature on the anti-*Trichomonas* activity of elderberry fruits and flowers. In this study, the extracts were prepared from fruits and flowers of *Sambucus* by using solvents with different polarities and are analyzed for their *in vitro* antiparasitic activity against *T. vaginalis*. While there have been recent studies on the antioxidant, antimicrobial, cytotoxic and antiviral activities of *S. nigra*^{6,8}, it is surprising that only one study has been conducted on the activity of *S. nigra* fruits against the common *T. vaginalis* parasite²⁴. This *in vitro* anti-*T. vaginalis* activity research will provide a basis for future *in vivo* trichomonocidal activity studies on *S. nigra*.

MATERIALS AND METHODS

Plant materials and preparation of extracts

S. nigra flowers and fruits were collected in June and August 2023, respectively. The plant species was authenticated by pharmacist Muammer Şen from Konya. The plant materials were dried at room temperature. The dried fruits and flowers were purchased from Pharmacist Muammer Şen, whose the owner of Temmuz Organic Farm, Selçuklu, Konya, Turkey. An amount of 5 g of fruits and flowers were grounded and macerated with solvent (25% hydroethanolic solution, 70% hydroethanolic solution, methanol and chloroform) with a plant/solvent ratio of 1:10 at room temperature under stirring for 48 hrs. The extracts were filtered through Whatman no.1 paper and solvent was evaporated to dryness by rotary evaporator under reduced pressure. The extracts were lyophilized and stored at -20°C until analysis²⁵.

Preparation of Trypticase Yeast Extract Maltose (TYM) Medium

Metronidazole resistant reference strain *T. vaginalis* ATCC 50143 was obtained from Celal Bayar University, Faculty of Medicine, Parasitology Bank, Manisa. In order to revive *T. vaginalis* parasites and to bring to logarithmic phase, trypticase yeast, maltose (TYM) medium was used. TYM medium is mainly

composed of 20 g trypticase, 5 g maltose, 10 g yeast extract, 0.2 g L-ascorbic acid, 1 g L-cysteine, 0.5 g agar and 0.8 g pH regulators (K_2HPO_4 - KH_2PO_4). Materials other than agar were weighed in the required amounts and dissolved in 900 ml distilled water using magnetic stirrer. The pH was adjusted to 6, followed by addition of agar and the mixture was sterilized for 20 minutes at 121°C under 15 psi. At the end of sterilization process, TYM medium was divided into 4 ml screw cap glass tubes. To make a total volume of 5 ml, before use, 100 IU/ml streptomycin, 100 IU/ml penicillin and commercially obtained inactive horse serum were added to TYM medium^{26,27}.

Revitalizing the parasites and bring into logarithmic phase

T. vaginalis strain was quickly thawed in a 37°C water bath within 1.5-2 minutes and then centrifuged and washed twice with fresh medium to remove remaining DMSO. After washing, the parasites accumulated at the bottom were added to TYM medium with 16% horse serum and incubated in the oven at 37°C. The growth of parasites was observed under a microscope. Then, the strains were then transferred to fresh medium to ensure average adaptation. Parasites that receive sufficient and necessary substances and pass the latent phase are expressed as production by multiplying their numbers in a generation period specific to their species is defined as logarithmic phase. Strains entering logarithmic phase were transferred to cell flask and incubated for the growth of parasites in high volumes^{27,28}.

Investigation of the antitrichomonal activity of extracts

Half maximal inhibitory concentration (IC₅₀) and minimum parasiticide concentration (MPC) values of the extracts against metronidazole-resistant *T. vaginalis* 50143 strain were determined *in vitro* by the liquid microdilution method in 96-well microplates²⁸. After the trophozoites entered the logarithmic phase, they were counted on the Thoma chamber using trypan blue stain and their viability was determined. In the experiments, trophozoites with a survival rate of over 95% were used and the amount of parasites was adjusted to 5×10^3 parasites/ml using a Thoma chamber. 5×10^3 parasites/ml *T. vaginalis* trophozoites were added to each well except for the negative control and the microplates were incubated at 37°C for 48 hrs. At the end of incubation, the motility of *T. vaginalis* trophozoites was evaluated by direct examination under an inverted microscope, and parasite viability was determined using trypan blue. For growth control (positive control), two wells containing *T. vaginalis* without tested extracts were used, and for contamination control (negative control), two wells containing the extracts without *T. vaginalis* were used. Then, the plates were incubated at 37°C under aerobic conditions. At the 24th and 48th hrs of incubation, the motility of *T. vaginalis* trophozoites was evaluated under an inverted microscope, and their viability was evaluated on a Thoma counting chamber using trypan blue stain. The extract concentration in the last well, where approximately half of the *T. vaginalis* trophozoites were determined to be alive using trypan

blue stain, was accepted as the IC₅₀ value and the MPC value was determined by taking trophozoites from the wells containing immobile parasites, passage them into a new TYM medium, and checking for growth at the 24th and 48th hrs²⁹.

Statistical analysis

The results of three parallel studies carried out at two different times were averaged and the viability percentages corresponding to all concentrations in which two-fold serial dilutions were analyzed with the Graphpad program and IC₅₀ and MPC values were expressed as µg/ml of the extracts.

RESULTS AND DISCUSSION

Table 1: Viability percentages of *T. vaginalis* trophozoites at 24 hrs in the presence of *S. nigra* fruits extracts.

Concentration (µg/ml)	70% hydroethanolic extract	25% hydroethanolic extract	Chlorofom extract	Methanol extract
25000	0.00±0.00	0.00±0.00	00.00±0.00	8.00±1.73
12500	0.00±0.00	40.33±0.57	0.00±0.00	50.00±0.00
6250	0.00±0.00	50.00±0.00	0.00±0.00	79.00±3.46
3125	13.33±2.88	63.33±0.57	10.66±2.30	92.33±2.88
1562.5	50.00±0.00	77.33±2.88	50.00±0.00	100.00±0.00
781.25	56.00±1.73	87.33±1.15	60.00±3.46	100.00±0.00
390.625	79.33±0.57	95.66±1.15	73.33±2.88	100.00±0.00
195.3125	95.66±1.15	100.00±0.001	93.66±23	100.00±0.00
97.65625	100±0.00	100±0.001	100±0.001	100.00±0.001

Table 2: Viability percentages of *T. vaginalis* trophozoites at 24 hrs in the presence of *S. nigra* flowers extracts.

Concentration (µg/ml)	70% hydroethanolic extract	25% hydroethanolic extract	Chlorofom extract	Methanol extract
25000	0.00±0.00	0.00±0.00	8.00±1.73	100±0.00
12500	0.00±0.00	0.00±0.00	50.00±0.00	100±0.00
6250	7.33±0.57	7.00±1.73	79.00±3.46	100±0.00
3125	50.00±0.00	50.00±0.00	92.33±2.88	100±0.00
1562.5	59.33±0.57	55.66±2.88	100.00±0.00	100±0.00
781.25	77.33±2.88	74.33±2.88	100.00±0.00	100±0.00
390.625	86.33±2.88	89.00±1.73	100.00±0.00	100±0.00
195.3125	99.00±1.73	100.00±0.00	100.00±0.00	100±0.00
97.65625	100±0.00	100.00±0.00	100.00±0.00	100±0.00

S. nigra fruit methanolic extracts' IC₅₀ were calculated as 1373 and 566 µg/ml for 24 hrs and 48 hrs respectively whereas no activity was observed at 48 hrs for the methanolic extracts of flowers even at concentration 25000 µg/ml. Among the flower extracts, the highest activity was observed in the chloroform extract with IC₅₀ value at 605 µg/ml. The 70% hydroethanolic extract of fruits exhibited the highest

In this study, the anti-*T. vaginalis* activity of *S. nigra* fruits and flowers extracts prepared with solvents of different polarities was comparatively analyzed. The parasite viability percentages of the fruit and flower extracts at 24 and 48 hrs are demonstrated in Table 1, Table 2, and Table 3, Table 4 respectively. The extracts of *S. nigra* reduced the number of live *T. vaginalis* parasites in a time and concentration dependent manner. The fruit extracts exhibited higher trichomonocidal activity than the flower extracts. As shown in Table 5, after 24 hrs of incubation the chloroform and 70% hydroethanolic extracts of fruits showed the highest inhibition of parasites with IC₅₀ values of 1136 and 1138 µg/ml respectively.

trichomonocidal activity with 513 µg/ml IC₅₀, while it was 8 µg/ml for metronidazole. In a study conducted on trichomonocidal activities of herbs used in Korea, the methanolic extracts of *Torilidis fructus*, *Sophora radix* and *Agrimoniae herba* were reported to show complete trichomonocidal activity at the concentration of 400 g/ml where as metronidazole used as positive control showed 0.94% viability at the concentration of 5 g/ml³³.

Table 3: Viability percentages of *T. vaginalis* trophozoites at 48 hrs in the presence of *S. nigra* fruits extracts.

Concentration (µg/ml)	70% hydroethanolic extract	25% hydroethanolic extract	Chlorofom extract	Methanol extract
25000	0.00±0.00	0.00±0.00	00.00±0.00	00.00±0.00
12500	0.00±0.00	0.00±0.00	0.00±0.00	00.00±0.00
6250	0.00±0.00	0.00±0.00	0.00±0.00	00.00±0.00
3125	0.00±0.00	8.66±0.57	0.00±0.00	00.00±0.00
1562.5	10.66±0.57	35.33±1.15	7.33±0.57	5.33±0.57
781.25	36.66±2.30	50.00±0.00	32.66±2.88	30.33±2.30
390.625	50.00±0.00	63.00±1.73	50.00±0.00	50.00±0.00
195.3125	74.66±0.57	77.00±1.73	60.00±1.73	66.66±5.77
97.65625	85.66±4.04	89.66±1.15	80±1.73	77.00±1.73

Table 4: Viability percentages of *T. vaginalis* trophozoites at 48 hrs in the presence of *S. nigra* flowers extracts.

Concentration (µg/ml)	70% hydroethanolic extract	25% hydroethanolic extract	Chlorofom extract	Methanol extract
25000	0.00±0.00	0.00±0.00	0.00±0.00	100±0.00
12500	0.00±0.00	0.00±0.00	7.33±1.15	100±0.00
6250	0.00±0.00	0.00±0.00	30.66±1.15	100±0.00
3125	5.66±0.57	7.33±1.15	50.00±0.00	100±0.00
1562.5	31.00±5.19	30.00±1.73	57.66±2.30	100±0.00
781.25	50.00±0.00	50.00±0.00	70.00±1.73	100±0.00
390.625	62.33±1.15	66.00±3.46	87.33±2.88	100±0.00
195.3125	78.66±0.57	76.00±1.73	97.00±1.73	100±0.00
97.65625	87.66±2.30	81.00±1.73	100.00±0.00	100±0.00

Table 5: The comparative *in vitro* activity of *S. nigra* fruit and flowers extracts on *T. vaginalis*.

Extracts	24. hrs (µg/ml)		48. hrs (µg/ml)	
	IC ₅₀	MPC	IC ₅₀	MPC
70% MM	1138	3125	513	1562
25% MM	8819	12500	886	3125
CMM	1136	3125	531	1562
MMM	1373	3125	566	1562
25% MC	2388	6250	1046	3125
70% MC	2611	6250	924	3125
CMC	2103	6250	605	2500
MMC	>12500	>12500	>12500	>12500
Metronidazole	14	66	8	33

MPC: Minimum Parasiticide Concentration; 70%:70% hydroethanolic extract; 25%: 25% hydroethanolic extract
 MM: extract of fruits; MC: extract of flowers. CMM: chloroform extract of fruits. CMC: chloroform extract of flowers
 MMM: methanol extracts of fruits; MMC: methanol extracts of flowers

In another survey on the anti-*T. vaginalis* activity, hydroalcoholic extracts of *Eugenia caryophyllata*, *Camellia sinensis* and *Terminalia chebula* were reported to have the best anti-trichomonal activity with IC₅₀ values of 1.21, 1.62 and 1.66 mg/ml, respectively³⁴. In the present study, the 70% hydroethanolic fruits extract which had the highest anti-*T. vaginalis* activity among the investigated fruit and flower extracts, inhibited the 50% of parasites at the concentration of 0.513 mg/ml which is much lower concentration than those extracts which were reported to have significant activity. In a previous report, methanolic extract of *S. nigra* fruits were evaluated for *in vitro* anti-*Trichomonas* activity at concentrations ranging from 100-800 µg/ml and at 800 µg/ml the extract was reported to have 100% efficacy after 48 hrs²⁴. In contrast, in this study fruit methanolic extract inhibited the 50% of parasites at 566 µg/ml at 48 hrs. 95% parasite inhibition was observed in methanol extract at 1562.6 µg/ml. Obtained results showed that *S. nigra* has acceptable efficacy *in vitro* and according to minimum parasiticide concentrations fruit extracts were significantly better than flower extracts.

However *in vivo* trichomonocidal activity search is required for further studies. Among the phytochemicals in *S. nigra*, flavonoids, anthocyanins, phenolic acids, cyanogenic glycoside and vitamin A and vitamin C have been reported⁸. Research has been conducted mostly on the phenolic components of the elderberry plant³⁰⁻³², but as observed in this study, chloroform extract, which is known to contain more lipophilic compounds, also showed strong activity. In addition to hydroethanolic and methanolic extracts of fruits, the chloroform extracts of fruits and flowers are also worth for investigating antitrichomonocidal candidates.

CONCLUSIONS

There are limited studies on the activity of elderberry against trichomoniasis. This study is the first comparative analysis of the anti-*T. vaginalis* activity of *S. nigra* extracts. This study will serve as a reference source in isolating the components of elderberry fruits with antiparasitic activity. Anti-*T. vaginalis* bioactivity guided assay is planned for further studies for the isolation of active constituents of fruits and flowers of *S. nigra*.

ACKNOWLEDGEMENTS

We would like to render our thanks to Parasitology Bank of Faculty of Medicine, Celal Bayar University for supplying the parasite strains.

AUTHOR'S CONTRIBUTION

Erdemir E and Çavus I: investigation, data curation, writing and editing. **Kayalar H:** plant material supply, methodology. **Kayalar H and Ozbilgin A:** supervision. Final manuscript was read and approved by all authors.

DATA AVAILABILITY

The data will be available to anyone upon request from the corresponding author.

CONFLICT OF INTEREST

None to declare.

REFERENCES

- Dündar S. Phtotherapeutic investigations on *Sambucus L.* species. Master Thesis 2009; Gazi University Health Institute.
- Taxonomy and Nomenclature; 2016. *Sambucus L.* Taxonomic serial no. 35315. Integrated taxonomic information system. http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_to_pic=TSN&search_value=35315
- Natural Resources Conservation Service, PLANTS database 2016. Classification for kingdom plantae down to genus *Sambucus L.* The United States Department of Agriculture. <http://plants.usda.gov/java/ClassificationServlet?source=display&classid=SAMBU>
- Kilham C. Health Benefits Boost Elderberry. Herbal Gram, American Botanical Council 2000; 50: 55.
- Charlebois D. Elderberry as a medicinal plant 2007; 284-292. J. Janick, and A. Whipkey (eds.), Issues in new crops and new uses. ASHS, Alexandria, VA.
- Jabbari M, Daneshfard B, Emtiazy M, Khiveh A, Hashempur MH. Biological effects and clinical applications of dwarf elder (*Sambucus ebulus L.*): A Review. J Evid Based Comp Altern Med 2017; 22(4):996-1001. <https://doi.org/10.1177/2156587217701322>
- The American Botanical Council. Clinical Guide to Elder Berry 2003.
- Młynarczyk K, Walkowiak-Tomczak D, Łysiak GP. Bioactive properties of *Sambucus nigra L.* as a functional ingredient for food and pharmaceutical industry. J Funct Foods 2018; 40:377-390. <https://doi.org/10.1016/j.jff.2017.11.025>
- Arslandoğlu Ş F, Sert S, Özdemir M. The importance of *Sambucus nigra* and *Sambucus ebulus*, which are widespread in Anatolian geography, as medicinal plants. Conference paper, Agromedya 2019; 58-62.
- Alıç B, Olcay N, Demir MK. Nutrient composition and functional properties of Black Elderberry (*Sambucus nigra L.*). JIST 2021; 11(2): 1140-1153. <https://doi.org/10.21597/jist.765296>
- Baytop, T. 1984. Türkiye'de Bitkiler ile Tedavi. İstanbul Üniversitesi Yayınları. No: 40, Sanal Matbaacılık. İstanbul.
- Kayabaşı N, Etikan S. The color obtained from the elderberry (*Sambucus nigra L.*) plant and the light and friction fastness of these colors on wool carpet yarns. J Agri Sci 1998; 4(3): 65-69. https://doi.org/10.1501/Tarimbil_0000000815
- Bergner P. Elderberry (*Sambucus nigra, canadensis*). Medical Herbalism, Winter 1996-97; 8(4): 11-12.
- World Health Organisation, Trichomoniasis, 2023. <https://www.who.int/news-room/factsheets/detail/trichomoniasis>
- Sena AC, Miller WC, Hobbs MM, et al. *Trichomonas vaginalis* infection in male sexual partners: Implications for diagnosis, treatment, and prevention. Clin Infect Dis 2007; 44:13–22. <https://doi.org/10.1086/511144>
- Cudmore SL, Delgaty KL, Hayward-McClelland SF, Petrin DP, Garber GE. Treatment of infections caused by metronidazole-resistant *Trichomonas vaginalis*. Clin Microbiol Rev 2004, 17(4):783-93. <https://doi.org/10.1128/CMR.17.4.783-793.2004>
- Kissinger P, Muzny CA, Mena LA, et al. Single-dose versus 7-day-dose metronidazole for the treatment of trichomoniasis in women: An open-label, randomised controlled trial. Lancet Infect Dis 2018; 18:1251–1259. [https://doi.org/10.1016/S1473-3099\(18\)30423-7](https://doi.org/10.1016/S1473-3099(18)30423-7)
- Smith JD, Cudmore SL, Garber GE. Diagnosis and treatment of metronidazole-resistant *Trichomonas vaginalis* infection. Antimicrob Drug Resist 2017, 1277-91. https://doi.org/10.1007/978-3-319-47266-9_28
- Wendel KA, Workowski KA. Trichomoniasis: challenges to appropriate management. Clin Infect Dis 2007;44(3): S123–S129. <https://doi.org/10.1086/511425>
- Hashemi N, Omidi D, Kheyri P, Khamesipour F, Setzer WN, Benchimol M. A review study on the anti-*Trichomonas* activities of medicinal plants. Int J Parasitol Drugs Drug Resist 2021; 15:92-104. <https://doi.org/10.1016/j.ijpdr.2021.03.001>
- Ziaei Hezarjaribi H, Nadeali N, Fakhari M, Soosaraei M. Medicinal plants with anti-*Trichomonas vaginalis* activity in Iran: A Systematic review. Iran J Parasitol 2019; 14(1):1-9.
- Niyati M, Joneidi Z, Kamalinejad M, Haghghi A, Valaei N, Abdi A. Anti- *Trichomonas* effect of *Rheum ribes* and *Foeniculum vulgare* extracts on *Trichomonas vaginalis* in vitro. J Islamic Iranian Trad Med 2015;6(3):198208.
- Fouladvand M, Barazesh A, Tahmasebi R, Mohammadi K, Khorami S. Lethal effect of various derivatives of curcumin on *Trichomonas vaginalis* in vitro. Iranian South Med J 2018;21(2):116-24.
- Niknam A, Esboei BR, Chabra A. Anti-*Trichomonas vaginalis* effect of methanolic extracts of *Sambucus nigra* in comparison with metronidazole. Jundishapur J Nat Pharm Prod 2020; 15(4): e65872. <https://doi.org/10.5812/jjppp.65872>
- Ozbilgin A, Durmuskahya C, Kayalar H, Ostan I. Assessment of *in vivo* antimalarial activities of some selected medicinal plants from Turkey. Parasitol Res 2014; 113(1):165-73. <https://doi.org/10.1007/s00436-013-0846-3>
- Aksoy Gökmen A, Kayalar H, Pektaş B, Kaya S. Investigation of *in vitro* anti-trichomoniasis effect of *Nigella sativa* oil against *Trichomonas vaginalis*. Izmir Kâtip Çelebi University Faculty Health Sci J 2018; 3(3): 7-10.
- Turan Faraşat V, Çavuş İ, Özbilgin A. The effects of different concentrations of horse serum on *Trichomonas vaginalis* Cryopreservation. Turk Microbiol Contemporary Record 2022; 52(2):103-108. <https://doi.org/10.54453/TMCD.2022.10337>
- Clinical and Laboratory Standards Institute (CLSI) Performance Standards for Antimicrobial Susceptibility Testing: Twenty-Fourth Informational Supplement. CLSI Document M100-S24 2014; Wayne, 34(1).
- Özel Y, Çavuş I, Ünlü G, Ozbilgin A. Investigation of the antitrichomonal activity of cinnamaldehyde, carvacrol and thymol and synergy with Metronidazole. Türkiye Parazitoloj Derg 2024, 48(2): 726-7. <https://doi.org/10.4274/tpd.galenos.2024.91855>
- Akduman G, Korkmaz Ş, Taşkın T, Güneş FE. Cytotoxicity of *Sambucus nigra L.* on cancer cell line and *in vitro* antioxidant properties. Clin Exp Health Sci 2023; 13: 896-901. <https://doi.org/10.33808/clinexphealthsci.1190301>
- Srinivas Goud N, Prasad G. Antioxidant, antimicrobial activity and total phenol and flavonoids analysis of *Sambucus nigra* (elderberry). Int J Curr Pharm 2020; 12: 1. <https://doi.org/10.22159/ijcpr.2020v12i1.36829>
- Haş IM, Teleky BE, Szabo K, et al. Bioactive potential of elderberry (*Sambucus nigra L.*): Antioxidant, antimicrobial activity, bioaccessibility and prebiotic potential. Molecules 2023, 30; 28(7):3099. <https://doi.org/10.3390/molecules28073099>
- Kim YC, Ryu JS, Kim HY, Choi KM, Kim HS, Park H. Trichomonocidal activity of herbal extracts used in Traditional Medicine in Korea. Korean J Oriental Physiol Pathol 2006; 20(1): 171-173.
- Jafari M, Amini-Khoei H, Cheshmpanam M, Abdizadeh R. A survey on the anti-*Trichomonas vaginalis* effect of the hydroethanolic extract of various medicinal plants *in vitro*. JSUMS 2023; 25(1):1-6. <https://doi.org/10.34172/jsums.2023.627>