



RESEARCH ARTICLE

CHEMICAL COMPOSITION AND ANTIMICROBIAL ACTIVITY OF THE ESSENTIAL OILS OF FOUR VARIETIES OF *LIPPIA MULTIFLORA* IN BENIN

GANDONOU Dossa Clément^{1*}, BAMBOLA Bouraïma², TOUKOUROU Habib³, GBAGUIDI Ahokannou Fernand², DANSOU Christian¹, AWEDE Bonaventure⁴, LALEYE Anatole⁵, AHISSOU Hyacinthe¹

¹Laboratory of Enzymology and Biochemistry of Proteins, Faculty of Science and Technology, University of Abomey-Calavi, 01BP: 188, Cotonou, Benin.

²Pharmacognosie Laboratory /Institute of Research and Experimentation in Traditional Medicine and Pharmacopoeia (IREMPT) / Benin Center for Scientific Research and Innovation (CBRSI)/Faculty of Science and Technology, University of Abomey-Calavi, 01 BP 06 Oganla Porto-novo, Benin.

³Laboratory of Organic Pharmaceutical Chemistry, School of Pharmacy, Faculty of Health Sciences, University of Abomey-Calavi, Fairground Campus, 01 BP: 188, Cotonou, Benin.

⁴Unit of Teaching and Research in Physiology Faculty of Health Sciences, University of Abomey-Calavi, Cotonou, Benin.

⁵Cellogenetics and Cell Biology Laboratory, Faculty of Health Sciences, University of Abomey-Calavi 01BP 188 Cotonou, Benin.

Article Info:

Abstract



Article History:

Received: 3 June 2019
 Reviewed: 14 July 2019
 Accepted: 27 August 2019
 Published: 15 September 2019

Cite this article:

Clément GD, Bouraïma B, Habib T, Fernand GA, Christian D, Bonaventure A, Hyacinthe LAA. Chemical composition and antimicrobial activity of the essential oils of four varieties of *Lippia multiflora* in Benin. Universal Journal of Pharmaceutical Research 2019; 4(4): 15-20. <https://doi.org/10.22270/ujpr.v4i4.293>

*Address for Correspondence:

GANDONOU Dossa Clément, Laboratory of Enzymology and Biochemistry of Proteins, Faculty of Science and Technology, University of Abomey-Calavi, 01BP: 188, Cotonou, Benin. E-mail: gandonoufils@gmail.com

Objective: Present study involves the study of the chemical composition of the essential oils extracted from the leaves by gas chromatography and mass spectrometry of *Lippia multiflora* harvested in the regions of Kétou, Savalou, Bohicon and Mono and tested by the well diffusion method against pathogenic microorganisms.

Methods: The essential oils studied are terpene compounds, aromatic compounds, aliphatic compounds and other natural substances. The inhibition zone diameters determined allowed us to evaluate their degree of germ sensitivity of the strains tested. Essential oils extracted from *Lippia multiflora* harvested in these areas have the most pronounced antimicrobial activity.

Results: In total, the essential oils tested have different and specifically a degree of sensitivity against *Staphylococcus aureus* ATCC 25923, *Escherichia coli* ATCC 25922, *Proteus mirabilis* A24974, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* except that harvested in the Savalou who does not have no degree of sensitivity on *Pseudomonas aeruginosa*.

Conclusion: This work paves the way for food preservation with extracted natural substances and also the formulation of natural antimicrobials for this fact.

Keywords: Antibacterial, Benin, chemical composition, *Lippia multiflora*, volatile extracts.

INTRODUCTION

For thousands of years, plants have been occupied a prominent place in the life of man. All known civilizations have used either wild or cultivated plants to provide for their basic needs: food, shelter, clothing and also for their medical needs. The use of medicinal plants has been evolved gradually with the changing needs of humans. Over the centuries, a first distinction has been made between edible and toxic plants. Subsequently, accumulated empirical knowledge has been enabled humans to take plants as the essential source of drugs. The study of this knowledge by modern science gradually reveals some secrets of

nature that allow man to continue his evolution. Until the beginning of the 20th century, almost all medicines were herbal. Even today, modern medicine is highly dependent on plants whose therapeutic virtues have been confirmed. Whatever parts and shapes plants use, they are extremely rich in complex chemical structures. The metabolism of plants contains thousands of different constituents, which belong to very different chemical families, such as alkaloids, phenols, flavonoids, terpenoids, and steroids. These secondary metabolites, extraordinarily diversified, are widely exploited in various fields: in the culinary field as dyes and flavors, in the medicinal field as antibiotics, antioxidants, drugs, etc., and in the agricultural field as

pesticides. These are secondary metabolites of aromatic plants containing in high proportion very active molecules with antibacterial, insecticidal, fungicidal, acaricidal and cytotoxic properties¹¹. Food-borne diseases are nowadays public health problems²⁴ notwithstanding all the modern innovations in the direction of the improvement and the performance of the techniques and the hygiene of the production of the food. It has been estimated that more than 30% of the populations in industrialized countries suffer from these diseases every year, and more particularly in the year 2000, two million deaths from diarrheal diseases worldwide¹⁰. Microbial activity was the primary mode of food deterioration and many microorganisms involved in these food-borne conditions most often contribute to the loss of quality and safety. In fact, essential oils (EO) are the most exploited thanks to their broad spectrum of known biological activities. Many volatile compounds of essential oils are nowadays common ingredients of pharmaceutical preparations such as pulegone, menthol and thymol¹⁹; α -pinene, a highly demanded ingredient in the international market, was used in the manufacture of vitamin E¹³. Numerous studies have made it possible, on the one hand, to systematically study essential oils extracted from aromatic plants commonly used in traditional pharmacopoeia in Benin⁶ and^{5,4}, and on the other hand to highlight the effectiveness of certain essential oils in the context of integrated protection against certain insect pests of cereal stocks¹⁷. However, prior to current work, and no scientific study in Benin has focused on testing the efficacy of volatile extracts of aromatic plants against pathogenic microorganisms and alterations identified in foods. Faced with the resistance of microorganisms to synthetic antibiotics, current study undertook to propose a cheaper and more credible alternative solution with aromatic plants used in traditional medicine. Choice of current study, following an ethnobotanical survey, focused on *L. multiflora*¹². The present study therefore aims to analyze the chemical compositions of the volatile extracts of the retained aromatic plants and to test their effectiveness against certain pathogenic and alteration microorganisms identified in staple foods in Benin. It was a question, specifically to determine the antibacterial activities of the extracts from the diameters of zones of inhibition around the wells vis-à-vis microorganisms tested. The search for these compounds was launched with the aim of discovering new effective therapies against diseases that have not been treated, and to reduce the use of synthetic products that are harmful to humans and their environment. Like antibiotics, antioxidants and pesticides. Despite attempts to list plants and their extracts, the number of species studied remains relatively low given the growing need for natural compounds of aromatic and/or medicinal interest.

MATERIALS AND METHODS

Plant material

It consists of plant material (*L. multiflora*), microbial strains (bacteria, yeast and mold), biological and

chemical diagnostic equipment. Survey in current study was carried out in 2015 in the main towns of the four departments of Benin which are chosen because of their demography and their varied geographical areas. Present study carried out the survey in the areas devoted exclusively to the marketing of medicinal plants in the markets of these towns. The concerned regions are: Plateau, Mono, Zou and Collines. Precisely, the survey was carried out in the towns of Kétou for the Plateau regions, Houéyogbé for Mono Bohicon and Djidja for Zou and Savalou for Collines. A specimen of the plant is deposited and identified by the national herbarium of the University of Abomey-Calavi (Benin)⁸. The plant materials used in present study were made of leaves or leaf stems of plant selected based on their frequency use against opportunistic infections and arterial hypertension. After harvesting the plants, their fresh samples were dried for ten to fourteen days in the dark in a constant-temperature cabinet (conditioned air).

Bacterial strains

The bacterial strains used consisted of 04 clinical strains (*Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*) and 03 reference strains (*Escherichia coli* ATCC 25922, *Staphylococcus aureus* ATCC 25923, and *Proteus mirabilis* A 24974).

Biological and chemical diagnostic equipment:

The equipment used in the laboratory consists of conventional glassware sterile single-use plates of 90mm diameter, a micropipette P100 Gilson brand and culture media. The culture media and reagents used in the microbiological diagnosis come from BIORAD, MERCK, OXOID, BASINGSTORE, BIOMERIEUX and DifCO laboratories. A 24-hour pure colony portion from the Mueller Hinton medium of each strain was emulsified in 5ml of physiological saline to obtain a turbidity of 0.5 on the Mc Farland scale. The chemical diagnostic equipment is a GC, a spectrometer, Clevenger for extraction of essential oils.

Methods

The survey was carried out using the method used in a previous study²². This study was performed with 20 herbal medicine doctors chosen from the agenda of 'Programme National de la Promotion de la Médecine Traditionnelle' of Benin Ministry of Health. It was carried out in the selected areas in presence of people who master the mother tongues of these regions and the plants. The tools of the survey were composed of a questionnaire and interviews. During the survey, questions are asked are related to:

- Used plants to treat the hypertension high blood pressure
- Vernacular names of the used plants.
- Parts of the used plants
- How to make the medicines, and dosage?
- Administration and the remedies

Questions were asked to traditional herbal medicine doctors and the over-sixties people with traditional knowledge. A selection from the most used species and the helped pharmaco-chemistry point of view after consulting the different data banks, to retain only one

species belonging to only one botanical family. This species was harvested from July to August 2015 in their natural habitat in Benin. The botanical determinations are carried out at 'Herbier National' of

University of *Abomey Calavi* (Benin). This harvested species presents either a similar use in the other traditional pharmacopoeias or other popular uses.

Table 1: Synoptic review of methods for detection, isolation and identification of microbial strains tested.

Microbial strains	Origin	Detection Method	Isolation	Identification
<i>Escherichia coli</i>	Tomato sauce	ISO 7251. NPP (Sept.1994)	Confirmation Kovacs Reagent Indole Test	API 20E
<i>Klebsiella pneumoniae</i>	Cow milk	ISO 4832 NPP (July 1991)	Isolation on EMB	API 20E
<i>Staphylococcus aureus</i>	Cooked com dough	SO 6888. Baird-Parker, Nutritive Agar (June 2003)	Coagulase test	API Staph

Extraction of oils and calculation of yields

About 200 g of plant material from each plant were each subjected to hydro distillation with a Clevenger type apparatus for 3 hours used in previous study^{4,15,23}. The essential oil thus obtained in an essencier was collected in a small dark bottle containing anhydrous sodium sulphate (Na₂SO₄) and stored at 4°C until use. The yield R was expressed relative to the fresh plant material and its calculation was as follows:

$$R = \frac{M (HE)}{M (MV)} \times 100$$

Where, R: yield; M (HE): mass of essential oil obtained; M (MV): Mass of extracted plant material.

Chemical composition of essential oils Analysis by gas chromatography

The oils were analyzed using a VARIAN CP.3380 gas chromatograph equipped with two capillary columns (apolar and polar) and connected to a Varian integrator (Model C-R4A^o);²³.

Table 2: Yields of essential oils of the studied plants.

<i>L.multiflora</i>	Part analyzed	Yield
Bohicon (Djidja)	Leaf	2.18±0.04
Savalou (Chetti)	Leaf	1.07±0.03
Kétou (<i>Ikpilnè</i>)	Leaf	0.58±0.02
Mono (<i>Houéyogbé</i>)	Leaf	1.70±0.04

Analysis by gas chromatography and mass spectrometry

Technical characteristics of the equipment

All essential oils were analyzed by GPC / MS, which results in the separation of the constituents of the mixture on a column mounted in series with a mass spectrometer. Each component leaving the column passes directly into the ionization field of the spectrometer. The molecule subjected to the electronic impact is fragmented. The mass spectra obtained for each compound are compared with data from the literature as well as those from the laboratory data base. CPG/MS couplings were performed on a Hewlett Packard-Quadruple type device (Model 5970). The chromatograph was equipped with a fused silica capillary column 30 m long and 0.25 mm inside diameter with a dimethylpolysiloxane (DB-1) type grafted phase 0.25 µm thick.

Evaluation of the antibacterial activity of essential oils Isolation and purification of microbial strains:

The clinical strains were isolated from stool samples at the National Public Health Laboratory of Benin.




Figure 1: Arrangement of the wells on the 90 mm diameter petri dish.

Sensitivity test by well diffusion

Each inoculum was seeded by swabbing onto petri dishes containing Mueller Hinton agar. Using the sterile pasteur pipette tip 6 mm diameter wells were dug. Then using a cone and a micropipette 50µl of each extract were deposited in the wells previously dug. A well containing sterile distilled water to serve as a negative control. The Petri dishes were left for 1 hour at room temperature for pre-diffusion of the substances, before being incubated at 37°C in an oven for 18 h. After the incubation period, the dishes were examined for measurements of the inhibition zones. The antibacterial activity of the extracts was determined from the diameters of inhibition zones around the wells as shown (Figure 1).

RESULTS AND DISCUSSION

Yields of the plants studied

The essential oil yields of plants of the family Verbenaceae, the yields of essential oil extracted from the leaves of *L. multiflora* (Kétou, Savalou, Mono and Bohicon) are relatively high and their yields are 0.58±0.02%, 1.07±0.03%, 1.70±0.04 and 2.18±0.04%, respectively.

Chemical composition of essential oils

Present work involves the study of the chemical compositions of essential oils according to its large botanical family of different regions¹⁵.

Verbenaceae: This family includes *L. multiflora* (from Kétou, Savalou, Bohicon and Mono). As shown in Table 3, the oils obtained do not have the same composition. There is a relatively large difference in composition in oils recovered from *L. multiflora* from Savalou, Djidja, Kétou and Mono. Indeed, the major components of *L. multiflora* essential oil of Savalou are: 1,8-cineol 50,6%, α -terpinol 13,8%, aromadendrene 6,2%, germacrene - D 2,2%, linalool 0.3%, Isoledene 0.2%, Pinocarvone 0.2%, (E) - 0.1%

nerolidol, 0.1% 6-ol camphene, α -cubebene 0.1%, terpinene-4-ol 0.3%, sabinene 14.4%, α -pinene 5.7%, myrcene 2.9%, β -pinene 1.9%, γ -terpinene 0, 3%, α -tujene 0.3%, δ -cadinene 0.2%). Those of Djidja are: linalool 91%, germacrene D 2.9%, α -caryophyllene 2.1%, allo-aromadendrene 1.9%, (E) -olidolol 0.5%, δ -cadinene 0.4%, trans- β -ocimene 0.3%, α -copaene 0.2%, 1,8-cineol 0.2%, α -muurolene 0.1%, Romadendrene 0.1%, caryophyllene oxide 0.1%, 0.1% sabinene, 0.1% myrcene.

Table 3: Chemical compounds of the oil extracted from the leaves of *L. multiflora* harvested in the area of Savalou, Bohicon, Mono and Kétou.

Chemical composition	Savalou (Tchetti)	Bohicon (Djidja)	Mono (Houéyogbé)	Kétou (Ikpilè)
camphèn-6-ol	0.1			
α -cubebène	0.1			
(E)- nerolidol	0.1	0.5		
Terpinolène	0.2			
Linalool	0.3	91		
Terpinen-4-ol	0.3		12,50	
β -pinène	1.9			
γ -terpinène	0.3			
α -thujène	0.3			
Pinocarvone	0.2			
δ -cadinene	0.2	0.4		
Isoledene	0.2			
Germacrene – D	2.2	2.9		4.84
Myrcène	2.9	0.1		
α -pinène	5.7		4.50	4.65
Aromadendrène	6.2			
α -terpinol	13.8		13	
Sabinène	14.4	0.1		
1,8-cinéol	50.6	0.2		13.50
Romadendrène		0.1		
Oxyde de caryophyllène		0.1	10.25	
α -muurolène		0.1		
α -copaène		0.2		
Trans- β -ocimène		0.3		
Allo-aromadendrène		1.9		
α -caryophyllène		2.1		
β -Myrcene			1.98	1,92
beta-Caryophyllene				28
gamma-Terpinène			13.65	47
Methanethiol			08.12	
Beta Pinene			08	
Carvacrol			13	
p-cymène			15	

Table 4: Standard used for reading the results of the antibiogram tests of plant extracts.

Inhibition zone diameter	Degree of sensitivity of the germ	Symbol
$\Delta < 7\text{mm}$	Insensible	-
$7\text{mm} \leq \Delta < 8\text{mm}$	Sensible	+
$8\text{mm} \leq \Delta < 9\text{mm}$	Assez sensible	++
$\Delta \geq 9\text{mm}$	Très sensible	+++

Those of Mono are: caryophyllene oxide 10.25%, Methanethiol 08.12%, carvacrol 13%, β -Myrcene 1.98%, α -pinene 4.5%, Beta Pinene 08%, p-cymene 15%, terpinen-4-ol 12.5%, α -terpinol 13%) and those of Kétou are: gamma-Terpinene 47%, beta-Caryophyllene 28%, 1,8-cineol 13.50%, Germacrene-D 4.84 %, α -pinene 4.65%, β -Myrcene 1.92%, α -Pinene 4.64%. These results explain many fluctuations in the values of the physicochemical properties of the various

essential oils obtained. Oxygen compounds represent respectively (99.8% for the plant of Djidia, 93.34% for the Kétou plant, 76.8% for the Savalou plant and 31.37% for the Mono plant) of the chemical composition essential oil of *L. multiflora*. The level of these compounds is relatively important in the essential oil Djidja, Kétou and Savalou, as well as in the other Mono with a respective difference \approx 68.43, 61.97 and 45.43%. Linalool was found in the oil of *L. multiflora*

of Djidja with a significant percentage (91%) than in *L. multiflora* of Savalou (0.3%), the difference is 90.7%. So study can concludes that the region of Djidja favors this compound in the set of essential oil.

Table 5: Inhibition zone diameters (mm) of the different bacterial strains.

Inhibition zone diameter (mm)				
Bacterial strains	LD	LS	LK	LM
<i>Staphylococcus aureus</i> ATCC 25923	40	34	32	30
<i>E. coli</i> ATCC 25922	30	29	30	27
<i>Proteus mirabilis</i> A24974	23	23	27	26
<i>Staphylococcus aureus</i>	34	32	00	33
<i>Escherichia coli</i>	30	27	30	60
<i>Klebsiella pneumoniae</i>	29	30	29	32
<i>Pseudomonas aeruginosa</i>	09	07	10	14

Same thing for the germacrene-D, (E)-nerolidol and δ -cadinene respective percentage (2.9%, 0.5% and 0.4%) found in the essential oil Djidja and Respective Aure (2, 2%, 0.1% and 0.2%) found in the essential oil of Savalou but with a very small difference of the order of (0.7%, 0.4% and 0.2%). Terpinen-4-ol was found in *L. multiflora* oil of Mono with a significant percentage (12.50%) than in *L. multiflora* from Savalou (0.3%), the difference was 12.2%. Myrcene is found in *L. multiflora* Kétou oil with a percentage (4.84%) than in *L. multiflora* Savalou (2.2%) whose difference was (2.64%). 1,8-Cineol was found in *L. multiflora* oil of Kétou with a percentage (13.50%) that in *L. multiflora* of Djidja (0,2%) whose difference was (13,3%). Caryophyllene oxide (10.25%) was found in the oil of *L. multiflora* of Mono and aure (0.1%) found in the essential oil of Djidja but with a difference of (10.15 %). Gamma-Terpinene was found in Ketu *L. multiflora* oil with a significant percentage (47%) as compared to Mono *L. multiflora* (13.65%), the difference being 33.35%. So it can we reported that the Kétou region favors this compound in the essential oil complex. Similar results were found with α -pinene. On the other hand note that some compounds are present in the essential oil of *L. multiflora* Savalou, with a much higher percentage than the essential oil of Djidja, Mono and Kétou with a more or less different rates. In this case, study can conclude the Savalou region of Djidja, Mono and Kétou has been a risk of having compounds well present but with very low percentages. For β -Myrcene (1.92-1.98%) and α -pinene (4.50-4.65%), it was found that they remained almost stable, did not undergo large changes. The variation detected in the chemical composition of the essential oil from the region of Savalou, Djidja and Aure Mono and Kétou, were related to several parameters such as: the environmental factor, the nature of the soil, climatic and geographical conditions that change from one region to another, and at the time of harvest²⁰. As for the total number of compounds found in *L. multiflora* Kétou was five compounds less than the other Savalou, Djidja and Mono which contain respectively nineteen, fourteen and ten compounds. The essential oil of *L. multiflora* is generally composed of molecules that the pharmaceutical industry often uses. Indeed, the

presence of α -pinene compounds, the most common molecule-marking terpene in nature, was an effective molecule for the treatment of respiratory conditions such as colds, coughs or bronchitis. Moreover, thanks to the similarity of its smell with that of fir, it was commonly used in deodorants and cleaning products. On the other hand, the presence of the compound (Z)- β -ocimene was known for its properties and its stabilities notably preventing its oxidation. This composition was different from that previously reported by^{1,8,14}.

CONCLUSIONS

The chemical composition of the essential oil extracted from the leaves of *L. multiflora* harvested in the regions of Kétou, Djidja, Savalou and Mono was studied by CPG and CPG coupled with mass spectrometry. Djidja oil is mainly composed of linalo, germacrene D and α -caryophyllene, the Savalou oil was made up of 1,8-cineol, sabinene, α -terpinol, the one of Mono composed of gamma-terpinene, -cymene, Terpinene, Caryophyllene oxide, Carvacrol, Methanethiol, Alpha-terpinene, Beta Pinene and Ketu Oils consist of sesquiterpenes including gamma-Terpinene, Cineole, beta-Caryophyllene, caryophyllene oxide. The different essential oils investigated are composed of molecules with terpene structures that act as isolation or synergistic effect of the majority and minority compounds they contain on the different bacteria. The different sensitivities determined on each of them, essential oils as well as their compositions (Table 3) act differentially on the isolated microbial strains and in the present study, it appears that the Gram + are more sensitive to their effect than the Gram. Through these works, the essential oil of *L. multiflora* offers hope. Indeed, this oil, commonly called "Gambia tea" is an aromatic plant very little known in Benin and used in the treatment of several pathologies in traditional medicine. Bibliographic research has also shown that this plant has several therapeutic virtues. The results of present work confirm these effects and show that this plant has excellent antimicrobial properties.

ACKNOWLEDGEMENTS

The authors sincerely thank all the actors of the laboratories involved in the success of this work. We also thank Mr. Richard FELICHETAND, Hyacinthe AHISSOU and Fernand Ahokannou GBAGUIDI for their technical assistance.

AUTHOR'S CONTRIBUTIONS

Clément GD: writing original draft, conceptualization. **Bouraima B:** methodology, investigation. **Habib T:** writing, review, resources. **Fernand GA:** writing, review, and editing. **Christian D:** methodology, investigation, formal analysis. **Bonaventure A:** writing, review, and editing, project administration. **Hyacinthe LAA:** conceptualization, methodology. Final manuscript was read and approved by all authors.

DATA AVAILABILITY

The datasets generated during this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

None to declare.

REFERENCES

- Agnani H, T Makani, A Akagah, C Menut, JM Bessière. Volatile constituents and antioxidant activity of essential oils from
- multiflora* Moldenke, growing in Gabon. *Flav. Fragr J* 2005; 20: 34-38. <https://doi.org/10.1002/ffj.1383>
- Ahoussi ED, Sessou P, Wotto VD, Yèhouenou B, Kinsoudé E, Kpatinvoh B, Soumanou M, Sohounhloué D. Development of a production technology of a local drink Africa drink based on essential oils of peppermint and green mints. *SOACHIM Newsletter*; 2010; 7: 39-53.
- Alitonou GA. Essential oils extracted from aromatic plants of Beninese origin. Chemical study, biological evaluation and potential applications. PhD thesis of the Universities of Abomey-Calavi and Montpellier II; 2006: 283. <https://doi.org/10.3390/medicines3040025>
- Avlessi F, G Alitonou, D Sohounhloué, C Menut, JM Bessière. Aromatic plants of Tropical West Africa. Part XIV, Chemical and biological investigation of *Lippia multiflora* Moldenke essential oil from Bénin. *J. Essential Oil Res* 2005; 17:405-407. <https://doi.org/10.1080/10412905.2005.9698944>
- Ayedoun MA, Sohounhloué DK, Menut C, Lamaty G, Bessière JM. Chemical composition of the essential oils of two species of *Cymbopogon* of Benin exploitable industrially. *African Bulletin Biores Energy Dev Env* 1997; 8: 4-6. <https://doi.org/10.1080/10496475.2016.1201718>
- Baba-Moussa F, Koumaglo K, Ayedou K, Akpagana K, Moudachirou M, Bouchet P. Antifungal activities of essential oils extracted in Benin and Togo. *Cryptogamy Mycol* 1997; 18: 165-168. <https://doi.org/10.1155/2017/8761610>
- Bassolé IHN, Lamien-Meda A, Bayala B, Tirogo S, Franz C, Novak J, Nebié RC, Dicko MH. Composition and antimicrobial activities of *lippia multiflora* moldenke, *mentha piperita* L. and *ocimum basilicum* L. Essential Oils and their major monoerpene alcohols alone and in combination. *Molecules* 2010; 15: 7825-7839. <https://doi.org/10.3390/molecules15117825>
- Bassolé IHN, AS Ouattara, R Nebié, CAT Ouattara, ZI Kaboré, SA Traoré. Chemical composition and antibacterial activities and *Lippia chevaleri* and *Lippia mutiflora* from Burkina Faso. *Phytochem* 2003; 62: 209-212. [https://doi.org/10.1016/S0031-9422\(02\)00477-6](https://doi.org/10.1016/S0031-9422(02)00477-6)
- Burt S. Essential oils: Their antibacterial properties and potential applications in foods-a review. *Int J Food Microbiol* 2004; 94: 223-253. <https://doi.org/10.1016/j.ijfoodmicro.2004.03.022>
- Faleiro L, GM Miguel, CA Guerrero, JMC Brito. Antimicrobial activity of essential oils of *Rosmarinus officinalis*, *Thymus mastichina* L. spp *Mastichina* and *Thymus albicans* in: Proceedings of the second WOCMAP congress on medicinal and aromatic plants. Part 2. Pharmacognosy, pharmacology, phytomedicine, toxicology Mendoza-Argentina; 1999 <https://doi.org/10.17660/ActaHortic.1999.501.4>
- Gandonou DC, Ahissou H, Tokoudagba JM, Dansou C. Ethnobotanical, phytochemical and toxicity analysis of a Beninese antihypertensive plant: *Lippia multiflora*. *Int J Biol Chem Sci* 2017; 11(4): 1816-1828. <https://doi.org/10.4314/ijbcs.v11i4.31>
- Ghanmi M. Chemical composition and antimicrobial activity of the turpentine species of maritime pine (*Pinus pinaster*) and *Aleppo pine* (*Pinushalepensis*) from Morocco. *Acta Bot Gallica* 2007; 154(2): 293-300.
- Gouollaly T, Nkounkou LC, Mahmoud Y, Ouamba JM, Abena AA, Chalchat JC, Figueiredo G. Variation in the chemical composition of the essential oils of different organs of domesticated *Lippia multiflora* Moldenke. *Afr J Biotech* 2010; 9(41): 7009-7013. <https://doi.org/10.5897/AJB10.344>
- Kamagaju L, Koumaglo K, Agbonon A, Nkurunziza JP, Kabera J, Senyazobe JMV. Chemical study of the essential oils of *Lippia multiflora* Moldenke and *Cymbopogon giganteus* chiov from Togo. *UNR-Rwandese Studies* 2006; 12:5-14. <https://doi.org/10.1080/10412905.2008.9699420>
- Menut C, Lamaty G, Sohounhloué DK, Dangou J, Bessière JM. Aromatic Plants of Tropical West Africa. III chemical composition of leaf essential oil of *Lippia multiflora* Moldenke from Benin. *J Essent Oil Res* 1995; 7. <https://doi.org/10.1080/10412905.2008.9699420>
- Noudogbessi JP. Chemical studies and biological activities of extracts of aromatic plants of Beninese origin. UAC single doctoral thesis and Université Blaise Pascal Clermont-Ferrand 2009; 329.
- OMS. World Health Organization (WHO) Traditional Medicine Report: Needs and Potential; 2002: No. 46.
- Pauli A. Antimicrobial properties of essential oil constituents. *Int J Aromather* 2001; 11: 126-133. <https://doi.org/10.3390/medicines3040025>
- Riyaha H. Thesis; valorisation of aromatic and medicinal plants: study of the chemical and antibacterial potential of essential oils of *Rosmarinus officinalis* (wild and domesticated); 2013.
- Talalaj S, Bissangou MF, Ouamba JM. Chemical valorization of some aromatic and medicinal species of Congo: *Ageratum conyzoides* L., *Chromolaena odorata* King and Robinson, *Hyptis suaveolens* Poit and *Lippia multiflora* Moldenke. *Pharm Med Trad Afr* 1997; 9: 70-80.
- Tokoudagba JM, Chabert P, Anger C, N'Gom S, Gbenou J, Moudachirou M, Schini-Kerth V, Lobstein A. Research of plants with antihypertensive potentialities in Beninese biodiversity. *Ethnopharmacologia* 2010; 44: 32-41. <https://doi.org/10.3389/fpls.2017.01845>
- Yehouenou BB. Chemical composition and antimicrobial activities of essential oils (EO) extracted from leaves of *Lippia rubosa* A. Chev against foods pathogenic and adulterated microorganisms. *African J Microbiol Res* 2002; 6(26): 5496-5505. <https://doi.org/10.5897/AJMR12.698>
- WHO. Food safety and Food-borne illness. World health organization Fact Sheet; 2002: 237 revised January Geneva.