

Available online at [www.ujpronline.com](http://www.ujpronline.com)

## Universal Journal of Pharmaceutical Research

An International Peer Reviewed Journal

ISSN: 2831-5235 (Print); 2456-8058 (Electronic)

Copyright©2019; The Author(s): This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited



### REVIEW ARTICLE

## ETHNOBOTANY, PHYTOCHEMISTRY AND PHARMACOLOGY OF *OCHNA SCHWEINFURTHIANA*: A REVIEW

Yusuf AJ<sup>1\*</sup> , Abdullahi MI<sup>1</sup> , Muhammad AA<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical and Medicinal Chemistry, Usmanu Danfodiyo University, Sokoto, Nigeria.

<sup>2</sup>Department of Pharmaceutical and Medicinal Chemistry, Ahmadu Bello University, Zaria, Nigeria.

#### Article Info:



#### Article History:

Received: 4 October 2019  
Reviewed: 7 November 2019  
Accepted: 28 December 2019  
Published: 15 January 2020

#### Cite this article:

Yusuf AJ, Abdullahi MI, Muhammad AA. Ethnobotany, phytochemistry and pharmacology of *Ochna schweinfurthiana*: A review. Universal Journal of Pharmaceutical Research 2019; 4(6):75-79.

<https://doi.org/10.22270/ujpr.v4i6.341>

#### \*Address for Correspondence:

Yusuf AJ, Department of Pharmaceutical and Medicinal Chemistry, Usmanu Danfodiyo University, Sokoto, Nigeria. Tel: 08036386793, E-mail: [amynajega@gmail.com](mailto:amynajega@gmail.com)

#### Abstract

*Ochna schweinfurthiana* (*Os*, Family: Ochnaceae) is a small evergreen tree used in ethnomedicine to treat different ailments; it is also used in agri-horticulture and as ornaments, dyes among others. It is a rich source of complex dimers of flavonoids and used for treatment of pain, inflammation, and arthritis. Chemical investigations carried out on the different parts of the plant have been confined to phenolic compounds majorly, bioflavonoids, glycosides, steroids and terpenes. The plant, *O. schweinfurthiana* have shown a wide spectrum of biological and pharmacological properties which include antimicrobial, cytotoxic/antiproliferative, genotoxicity, antinociceptive, anti-inflammatory, antioxidant and antiplasmodial. This review comprehensively summarizes the potential effects of the plant *O. schweinfurthiana*, chemically and pharmacologically. However, more researches in the aspect of phytochemical and biological studies are needed to exhaustively isolate bioactive compounds and evaluate their effects on other ailments as claimed by the traditional healers.

**Keywords:** Anti-microbial, antimalarial, flavonoids, *Ochna schweinfurthiana*, Ochnaceae, phenolics, pharmacological.

### INTRODUCTION

*Ochna schweinfurthiana* (*Os*) belonging to the Ochnaceae family is a small tree that was named after a German botanical collector and taxonomist Dr. Georg August Schweinfurth; it is a small tree that measures up to 4 m tall; the plant is commonly known as the brick-red *Ochna* in English, Jan-taru in Hausa language, Hiéké in Yoruba and Sa'aboule in Fofouldé<sup>1,2</sup>. The plant has found use as medicine, for agricultural, social and religious purposes<sup>1</sup>. This review centered on the ethnobotanical, phytochemical and pharmacological properties of *O. schweinfurthiana*.

#### Botanical Description

*Ochna* originated from a Greek word "*Ochne*" which means wild pear and it was named by Linnaeus in 1951 as *Ochna* because of the resemblance of their leaves with those of wild pear<sup>3</sup>. It is an old world genus of mainly trees, shrubs and shrublets which comprises of about 85 species<sup>4</sup> and it is widely distributed in tropical Asia, Africa and America<sup>5</sup> of which eleven (11) species are found in India<sup>6</sup>. *Ochna*'s are usually called Mickey Mouse plants, because of the appearance of the black druplets fruits. The Ochnaceae family is mainly

composed of trees and shrubs with an estimated 33 genera and 550 species<sup>7</sup> well distributed around the world especially in tropical Africa, Australia, Madagascar Asia, the Mascarene Island and America<sup>8</sup>. They are notably known for their unusual shiny leaves, with parallel veins that are closely spaced, alongside toothed margins with conspicuous stipules<sup>1,7</sup>. The largest genera are *Ouratea*, *Ochna*, *Campy lospermum*, *Sauvagesia* and *Quiina* with (200, 85, 65, 39 and 34 species) respectively (Table 1).

#### Morphology

*O. schweinfurthiana* is a small evergreen tree or shrub that grows up to 4 m tall and it has a dark grey bark that is fissured and cracked, separating into square segments<sup>10</sup>. The leaves are olive-green (1-13.5x1.7-5.5 cm) that oblanceolate to oblong or elliptic, apex somewhat rounded, base tapers into the petiole, margins rather bluntly toothed (serrulate), sometimes appearing almost scalloped, net-veining conspicuous on the upper surface and young leaves are coppery<sup>1,11</sup>. It bears bright yellow flowers (1.5 cm diameter) which are sweetly-scented from September to November, very short-lived, normally appearing before or with the young leaves<sup>10</sup>. In addition, it appears in a condensed

receme with 4–10 flowers on a short central stem and the petals fall very early<sup>11</sup>.

**Table 1: Ochnaceae subfamilies and their estimated number of species.**

Subfamily	Estimated number of species
Ouratea	200
Ochna	85
Campylospermum	65
Sauvagesia	39
Quiina	34
Total	423



**Figure 1: Leaf and fruits of *O. Schweinfurthiana*.**

The fruits of *O. schweinfurthiana* are 1-5 oval appear between August and January, attached at the base are 2-4 black berries when ripe; they are enlarged, borne on brick-red persistent sepals turning cherry to brick-red. The bark is dark grey, thick, and deeply fissured into a grid-like pattern<sup>10,11</sup>.



**Figure 2: Whole plant of *O. Schweinfurthiana*.**

#### Taxonomy

Kingdom: Plantae                      Order: Malpighiales  
 Family: Ochnaceae                      Genus: Ochna  
 Subfamily: Ochnoideae  
 Species: *O. schweinfurthiana*<sup>9</sup>

#### Common names

English: Brick-red Ochna, Hausa: Jan-taru  
 Yoruba: Hiéké, Fulfulde: Sa'aboule

#### Habitat, Distribution and Ecology

The plant grows in open deciduous woodland in tropical regions in Africa from Guinea to southern and northern Nigeria and across central Africa to Sudan, Uganda, Zimbabwe, Mozambique, Tanzania and Angola. It has medium water requirement when young and grows fast, flowers from September to

November. It required low maintenance and attracts insects and birds<sup>4,10</sup>.

#### Ethnomedicinal uses

Several preparations (powdered and decoctions) of the leaves and/or root of the *O. schweinfurthiana* have found general use as antimicrobial (wound dressing, eye infection), analgesic, anti-inflammatory and anthelmintic agents<sup>1</sup>. The leaf is also used as an antiseptic, stimulant, febrifuge, laxative, enema etc<sup>1</sup>. In Northern Cameroon, *O. schweinfurthiana* is used to treat different diseases such as rubella, burns, stomachache and multiple sclerosis<sup>12</sup>; the root of the plant is also used in the treatment of stomach and eye aches as well as headache, while the leaves are used for toothaches treatment<sup>2</sup>.

The pulverized bark is used to treat malaria, febrifuges and as anthelmintic, while the decocted root leaves and/or bark is used in wound dressing<sup>12</sup>. In Northern Nigeria, the *O. schweinfurthiana* is used to treat typhoid fever, measles and fungal skin infections<sup>12</sup>. The macerated roots of *O. schweinfurthiana* has been reportedly used to induce/speed the delivery process and for miscarriage<sup>13</sup>.

#### Other uses


The plant is used in agri-horticulture; the bark and flowers *O. schweinfurthiana* are cultivated for ornaments, dyes, stains, inks, tattoos and mordant among others. The wood is used for farming, forestry, hunting and fishing apparatus. The leaf has social, religious, superstitious and magic values among others<sup>1</sup>.

#### Phytochemical screening

Abdullahi et al.,<sup>12</sup> reported the presence of flavonoids, steroids/terpenes and saponins in the acetone leaf extract of *O. schweinfurthiana* and the methanol leaf extract indicated the presence of flavonoids and saponins. However, flavonoids, saponins, glycosides, tannins and steroids/terpenes were reported on the methanol stem extract of *O. schweinfurthiana*<sup>14</sup>. A study conducted by Ibrahim et al., reported the presence of carbohydrates, steroids/triterpenes, glycosides, saponins, tannins and flavonoids in the methanol leaf extract of *O. schweinfurthiana*<sup>15</sup>.

#### Bioactive constituents

Chemical constituents isolated from *O. schweinfurthiana* fall under the following class of secondary metabolites phenolics-flavonoids, and glycosides. *O. schweinfurthiana* have been reported to contain phenolic derivatives (such as flavonoids, bioflavonoids) which appear as free or in polymerized forms. Isolation and characterization of quercetin-3-O-β-D-glucopyranosyl-(1→6)-α-rhamnoside (quercetin rutinoside) from the *n*-butanol soluble fraction of methanolic leaf extract of *O. schweinfurthiana* was reported<sup>16</sup>. A novel biflavonoid compound, tri-methoxy lophirone was isolated from the chloroform soluble fraction of the methanol root extract of *O. schweinfurthiana*<sup>17</sup>.



**Figure 3: Chemical constituents isolated from *O. schweinfurthiana*.**

Ndongo *et al.*,<sup>18</sup> reported the isolation of seven flavonoids, hemerocallone, 6,7-dimethoxy-3',4'-dimethoxyisoflavone, amentoflavone, agathisflavone, cupressuflavone, robustaflavone, and epicatechin, and three other chemical constituents, lithospermoside, β-D-fructofuranosyl-α-D-glucopyranoside and 3β-O-D-glucopyranosyl-β-sitosterol from the ethyl acetate the stem bark extract of *O. schweinfurthiana*.

The roots of *O. schweinfurthiana* were reported to contain three new compounds viz; 4'''-methoxylophirone A, 4,4',4'''-trimethoxylophirone A. In addition, six known compounds were also isolated and characterized, including calodenone, calodenine B, lophirone A, gerontoisoflavone A, and 3β-O-D-glucopyranosyl-β-sitosterol. Six known compounds

were isolated from the powdered bark of *O. schweinfurthiana* and they include, hemerocallone, 6,7-dimethoxy-3',4'-dimethoxyisoflavone, lithospermoside, amentoflavone, agathisflavone and β-D-fructofuranosyl-α-D-glucopyranoside<sup>19</sup>.

#### **Biological and pharmacological activities**

##### **Antimicrobial activity**

The acetone and methanol leaf extracts of *O. schweinfurthiana* had a remarkable antibacterial effect against *S. aureus*, *S. typhi*, *K. pneumonia* and *P. aeruginosa* with a zone of inhibition ranging from 15 – 21 mm; the extracts had an MIC and MBC values of 10–20 mg/ml and 20–40 mg/ml, respectively<sup>12</sup>. Quercetin-3-O-β-D-glucopyranosyl(1→6)-α-rhamnoside from the *n*-butanol soluble fraction of the

methanolic leaf extract of *O. schweinfurthiana* showed an *in vitro* inhibitory effect against some bacterial isolates such as *S. aureus*, *MRSA*, *S. pyogenes*, *E. coli*, *K. pneumoniae*, *S. typhi* and *P. aeruginosa* with an MIC and MBC range between 2.5–5.0 and 5–20 µg/ml, but there was no effect against *B. subtilis*, *C. ulcerans* and *C. albicans*<sup>16</sup>. Earlier studies showed that trimethoxy lophirone A from the chloroform soluble fraction of the methanol root extract of *O. schweinfurthiana* inhibited the growth of some selected human pathogens including *S. aureus*, *S. pyogenes*, *P. aeruginosa*, *K. pneumoniae* and *S. typhi* with an MIC and MFC values of 5 µg/ml and 20 µg/ml, respectively<sup>17</sup>. Crude methanol stem extract of *O. schweinfurthiana* and its chloroform (CF), ethylacetate (EF) and *n*-butanol (BF) fractions inhibited the growth of *MRSA*, *S. aureus*, *S. pyogenes*, *S. typhi*, *S. dysenteriae*, *K. pneumoniae*, *N. gonorrhoea*, *P. aeruginosa*, *C. albicans*, *C. tropicalis*; the mean zone of inhibition of extract and fractions ranges from 20–29 mm; moreover, chloroform fraction showed greater antimicrobial activity with an MIC value of 1.25 mg/ml against all the test organisms except *P. aeruginosa*<sup>14</sup>.

#### Cytotoxic and antiproliferative effect

The methanol and ethylacetate stem bark extracts of *O. schweinfurthiana* demonstrated cytotoxicity against HeLa cells; amentoflavone and agathisflavone were also active<sup>18</sup>. Antiproliferative effect of *O. schweinfurthiana* extract was evaluated against *Glioblastoma multiforme* (GBM U-1242 MG) cell line and the extract reduced cell count by 20% with an IC<sub>50</sub> value 823.51 µg/ml<sup>20</sup>. The aqueous stem bark of *O. schweinfurthiana* did not show any cytotoxic effect on Vero monkey kidney cell line after 48 h incubation with an LC<sub>50</sub> value 50±1 µg/ml<sup>19</sup>.

#### Genotoxicity

Djova et al., reported that the extracts of *O. schweinfurthiana* were non genotoxic in a study they carried out; as none of the plant extracts demonstrated a dose dependent increase or revertent colonies ≥ the number of negative control revertent colonies; thus, the plant *O. schweinfurthiana* may not contain any genotoxic substances that may lead to mutations either by substitution or by reversion in the genetic material<sup>19</sup>.

**Antinoceptive and anti-inflammatory effect:** The methanol leaf extract of *O. schweinfurthiana* significantly inhibited the writhing response induced by acetic acid in a dose dependent manner; the highest dose exhibited maximum inhibition of pain (84.3%). In addition, the extract was also able to attenuate pain response in a similar manner though with a slower onset of action in the tail flick model<sup>15</sup>. The aqueous bark extract of *O. schweinfurthiana* exhibited good anti-inflammatory effect in both ferrous oxidation-Xylenol Orange (Fox) and BSA denaturation assays; the extract demonstrated good 15-lipoxygenase inhibitory effect with an IC<sub>50</sub> value of 32.2±0.36 µg/ml, however, an IC<sub>50</sub> of 130±5.78 µg/ml was recorded by the extract in the inhibition of heat induced albumin denaturation<sup>19</sup>.

#### Antioxidant effect

Messiet et al., evaluated the antioxidant activity of some compounds including 4"-methoxylophirone A,

calodenone, calodenine B, lophirone A, gerontoisoflavone A from the roots of *O. schweinfurthiana* using DPPH radical scavenging and ferric reducing-antioxidant power (FRAP) assays<sup>2</sup>. In the DPPH radical scavenging assay, calodenine B showed prominent effect with SC<sub>50</sub>=0.17±0.04 µM and EC<sub>50</sub>=4.25 µM, gerontoisoflavone A exhibited weak activity in all the models applied with SC<sub>50</sub>=19.00 µM and SC<sub>50</sub>=78.67 µg EAA/mg/dw in DPPH and FRAP respectively. The antioxidant property of the leaf, stem bark and fraction of *O. schweinfurthiana* was evaluated<sup>21</sup>.

#### Antiplasmodial effect

An *in vivo* study showed that the methanol leaf extract of *O. schweinfurthiana* exerted a suppressive effect against *Plasmodium berghei* at a lower dose (50 mg/kg); Ibrahim et al., concluded that, the extract possess blood schizonticidal activity as it was able to suppress malaria at the early stage<sup>22</sup>. Moreso, the extract reduced the level of parasitaemia with 100% cure at the lowest dose (50 mg/kg); the percentage inhibition of parasitaemia was higher than the chemo suppression which might be due to non-selectivity of the extract to the proliferative process of the parasite<sup>22,23</sup>.

Antiplasmodial effect of the ethylacetate roots extract of *O. schweinfurthiana* and some compounds including 4"-methoxylophirone A, (4E,7Z)-3,8-dicarboxy-1-(*O*-β-D-glucopyranosyl-(1→6)-*O*-β-D-glucopyranosyl-2,9 dihydroxyhexeicosa-4,7-diene, calodenine B, lophirone A and gerontoisoflavone A were investigated *in vitro*; 4"-methoxylophirone A showed good antiplasmodial effect against *P. falciparum* strain 3D7; this effect as explained by Messi et al.,<sup>2</sup> might be related to the presence of a methoxy group on position C-4" which has been known to enhance lipophilicity thereby enhancing its movement into the cells<sup>24</sup>; other compounds were found to be inactive<sup>2</sup>. Cold and hot aqueous leaf extracts of *O. schweinfurthiana* possess inhibitory effect against *P. falciparum in vitro*; thus there was significant reduction of parasitaemia.

The high dose (80 µg/ml) exhibited 86.42% (cold extract) and 85.06% (hot extract) reduction of parasitaemia. On the other hand, no significant difference was observed on the plasmodium lactate dehydrogenase (PLDH) activity of the treated extract when compared with the standard drug<sup>25</sup>.

#### Toxicity

Toxicity level of *O. schweinfurthiana* was assessed in mice both orally and intraperitoneally. The methanol leaf extract of the plant produced an LD<sub>50</sub> 774.6 mg/kg, i.p. while the oral LD<sub>50</sub> value for the extract was about 5000 mg/kg; according to this study, the leaf of *O. schweinfurthiana* is intraperitoneal toxic and orally safe<sup>22</sup>.

## CONCLUSIONS

*O. schweinfurthiana* exhibit a variety of biological effects; the plant is considered to be effective against cancer, malaria, oxidative stress, pain, inflammation and a wide range of activity against microbes; thus, the

pharmacological actions have been attributed to the presence of different classes of secondary metabolites such as biflavonoids, glycosides, steroids and terpenes among others. In addition, the mechanisms of action of the observed effects and evaluation of other pharmacological properties of *O. schweinfurthiana* need closer attention and it should be the objective of new researches on *O. schweinfurthiana*.

## ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Usmanu Danfodiyo University, Sokoto, Nigeria to provide necessary facilities for this work.

## AUTHOR'S CONTRIBUTION

**Yusuf AJ:** writing original draft, methodology. **Abdullahi MI:** investigation, formal analysis, conceptualization. **Muhammad AA:** writing, review and editing. Final manuscript was read and approved by all authors.

## DATA AVAILABILITY

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## CONFLICT OF INTEREST

No conflict of interest associated with this work.

## REFERENCES

- Burkill MH. The useful plants of west Tropical Africa, Families J–L. Royal Botanic Gardens Kew 1985; 4: 275.
- Messi AN, Mbing JN, Ndongo JT, et al. Phenolic compounds from the roots of *Ochna schweinfurthiana* and their antioxidant and antiparasitic activities. *Phytochemistry Lett* 2016; 17: 119-125. <https://doi.org/10.1016/j.phytol.2016.07.011>
- Muema MJ. Phytochemical and antimicrobial investigation of *Ochnathomasiana* Engl. and Gilg. MSc Thesis, Department of Chemistry, Kenyatta University 2015; 30– 35. [http://dx.doi.org/10.1093/jac/48.suppl\\_1.5](http://dx.doi.org/10.1093/jac/48.suppl_1.5)
- Verdcourt B. Ochnaceae. Flora of tropical east Africa. Royal Botanic Gardens, Kew, Richmond, United Kingdom 2005; 60.
- Rendle AB. The classification of flowering plants. Cambridge University Press, London. 1952; 2:12-18.
- Kirtikar KR, Bussu BD. Indian medicinal plants. Periodical expert book agency, Delhi. 2012; 2:1539.
- Christenhust MJM, Byng JW. The number of known plants species in the world and its annual increase. *Phytotexa* 2016; 261(3): 201-217.
- Mabberley DJ. Mabberley's plant-book. A portable dictionary of plants, their classification and uses. Cambridge university press, Cambridge 2008; 342.
- Simpson DP. Cassell's Latin Dictionary (5 ed.). London: Cassell Ltd. 1979; 883. ISBN 0-304-52257-0.
- Hyde MA, Wursten BT, Ballings P, Coates PM. Flora of Zimbabwe: Species information: *Ochna schweinfurthiana* F. Hoffm. [www.zimbabweflora.co.zw/speciesdata/species.php?speciesid.2019](http://www.zimbabweflora.co.zw/speciesdata/species.php?speciesid.2019); retrieved on 28 September, 2019.
- Anonymous. *Ochna schweinfurthiana*: Plant Book. Available online at [www.plantbook.co.za/ochna-schweinfurthiana](http://www.plantbook.co.za/ochna-schweinfurthiana) 2019; retrieved 28 September, 2019
- Abdullahi MI, Iliya I, Haruna AK, Sule MI, Musa AM, Abdullahi SM. Preliminary phytochemical and antimicrobial investigations of leaf extracts of *Ochna schweinfurthiana*. *African J Pharm Pharmacol* 2010; 4(2): 083-086.
- Bruschi P, Morganti M, Mancini M, Signorini MA. Traditional healers and laypeople: A qualitative and quantitative approach to local knowledge on medicinal plants in Muda (Mozambique). *J Ethnopharm* 2011; 138: 543-563. <https://doi.org/10.1016/j.jep.2011.09.055>
- Danmusa UM, Nasir IA, Abdullahi MI, Ahmad AA, Abdulkadir IS. Phytochemical analysis and antimicrobial activities of methanolic stem extracts of *Ochna schweinfurthiana* F. Hoffm. *J Pharmacy Pharmacog* 2015; 3(6): 171-182.
- Ibrahim ZYY, Musa AM, Abdullahi MI, Uba A, Yusuf AJ, Aliyu IM, Ya'u J. Phytochemical and antinociceptive studies on the *Ochna schweinfurthiana*(Ochnaceae). *Int J Advances Pharm Biol Chem* 2015a; 4(4): 838-843.
- Abdullahi MI, Musa AM, Haruna AK, et al. Anti-microbial flavonoid diglycoside from the leaf of *Ochna schweinfurthiana* Hoffm. (Ochnaceae). *Nigerian J Pharm Sci* 2011; 10(2): 1-7. <https://doi.org/10.1016/j.phytol.2016.07.011>
- Abdullahi MI, Musa AM, et al. Isolation and characterization of anti-microbial biflavonoid from the chloroform-soluble fraction of methanol root extract of *Ochna schweinfurthiana* (Ochnaceae). *African J Pharm Pharmacol* 2014; 8(4): 93-99. <https://doi.org/10.1076/phbi.41.2.92.14245>
- Ndongo JT, Issa ME, Messi AN, et al. Cytotoxic flavonoids and other constituents from the stem bark of *Ochna schweinfurthiana*. *Natural Prod Res* 2015; 17:1684-1687. <https://doi.org/10.1080/14786419.2014.991321>
- Djova SV, Nyegue MA, Messi AN, Afagnigni AD, Etoa FX. Phytochemical study of aqueous extract of *Ochna schweinfurthiana* F. Hoffm powder bark and evaluation of their anti-inflammatory, cytotoxic and genotoxic properties. *Hindawi Evidence-Based Complementary and Alternative Medicine*, 2019. <https://doi.org/10.1155/2019/8908343>
- Abdullahi MI, Musa AM, Tajuddeen N, Mohammed GM, Yusuf AJ, Iliya I. Anti-proliferative study and isolation of Ochnaflavone from the ethylacetate-soluble fraction of *Ochnakibbiensis* Hutch and Dalziel. *Nat Prod Res* 2016; <https://doi.org/10.1080/14786419.2016.1274892>
- Nyegue MA, Ngo Mbing J, Djova SV, et al. Evaluation of the antioxidant activity of the leaves, stem-barks extracts and fractions of *Ochna schweinfurthiana* F. Hoffm (Ochnaceae). *African J Pharm Pharmacol* 2016; 10(17): 370–378. <https://doi.org/10.1155/2019/8908343>
- Ibrahim ZYY, Abdullahi MI, Musa AM, Maje IM, Yusuf AJ, Aliyu IM. Acute toxicity profile and *Plasmodium berghei* inhibitory activity of *Ochna schweinfurthiana* (F. Hoffm) Ochnaceae leaf extract in Laboratory animals. *Int J Pharma Sci Res* 2015b; 6(10):1302–1306.
- Salawu OA, Tijani AY, et al. Anti-malarial activity of ethanolic stem bark extract of *F. bialbida* (Del) a. Chev (Mimosoidae) in mice. *Arc App Sci Res* 2010; 2: 261-268. <https://doi.org/10.1155/2011/618209>
- Monks MR, Ferraz A, Bordignon SA, et al. *In vitro* cytotoxicity of extracts from Brazilian Asteraceae. *Pharm Biol* 2002; 40: 494-500. <https://doi.org/10.1076/phbi.40.7.494.14681>
- Omoniwa BP. *In-vitro* antiparasitic activity of cold and hot aqueous extracts of *Ochna schweinfurthiana* leaf. *J Bacteriol Parasitol* 2017; 8(5 Suppl): 30.