



RESEARCH ARTICLE

EPIDEMIOLOGY, BACTERIAL PROFILE, AND ANTIBIOTIC SENSITIVITY OF LOWER RESPIRATORY TRACT INFECTIONS IN SANA'A AND DHAMAR CITY, YEMEN

Abdulrahman Y. Al-Haifi¹ , Abdul Salam Mohamed Al Makdad² , Mohammed Kassim Salah² , Hassan A. Al-Shamahy^{2*} , Wadee Abdullah Abdulwahid Al Shehari³

¹Department of Microbiology, Faculty of Medicine, Dhamar University, Dhamar, Yemen.

²Department of of Medicine, Faculty of Medicine, Dhamar University, Dhamar, Yemen.

³Medical Microbiology and Clinical Immunology Department, Faculty of Medicine and Health Sciences, Sana'a University, Republic of Yemen.

Article Info:



Article History:

Received: 2 February 2020

Reviewed: 10 March 2020

Accepted: 26 April 2020

Published: 15 May 2020

Cite this article:

Al-Haifi YA, Al Makdad ASM, Salah MK, Al-Shamahy HA, Al Shehari WAA. Epidemiology, bacterial profile, and antibiotic sensitivity of lower respiratory tract infections in sana'a and Dhamar city, Yemen. Universal Journal of Pharmaceutical Research 2020; 5(2):32-37.
<https://doi.org/10.22270/ujpr.v5i2.386>

*Address for Correspondence:

Dr. Hassan A. Al-Shamahy, Faculty of Dentistry, Sana'a University, P.O. Box 775 Sana'a, Yemen, Tel: +967-1-239551.
 E-mail: shmahe@yemen.net.ye

Abstract

Objectives: Lower respiratory infections (LRTIs) are the leading reason of death infectious diseases in the world and the fifth leading cause of death in general. The study aimed to identify the general characteristics of LRTI, the causative bacteria and the results of sensitivity to antibiotics.

Subjects and methods: A multicentre prospective study was performed at 3 University hospitals. The study included 555 clinical diagnostic cases as LRTI cases, 328 male and 227 female, aged 3 to 69 years. Clinical and demographic data were collected in the standard questionnaire, and samples included sputum or bronchial lavage (BAL) staining and culture. Samples were cultured in 3 different bacterial media, blood agar and LJ slope, chocolate agar with CO₂; cultures were then examined for possible bacterial pathogens of LRTI. Possible bacterial pathogens were isolated and identified by standard laboratory techniques, and microbial sensitivity testing was carried out by disc diffusion method.

Results: LRTI was recorded among all age groups and with less frequency in children less than 16 years of age. A large number of LRTI (36.2%) was not diagnosed, most in CAP (52.4%), followed by HAP (33.9%) while unidentified cases were lower in AECOPD (22.8%). CAP isolates are *K. pneumoniae* (26.2%), *S. pyogens* (12.3%), and *S. pneumoniae* (9%); in HAP are MSSA (24%), *E. Coli* (12.9%), MRAS (11.1%), *K. pneumoniae* (10.5%) and *P. aeruginosa* (7%); and in AECOPD are *M. catarrhalis* (47.2%), *K. pneumoniae* (17.2%), *H. influenzae* (10.7%) and *P. aeruginosa* (2%). In Gram-positive bacteria, high resistance to ampicillin/sulbactam (100%) and amoxicillin/clavulanate (100%) was recorded, while moderate resistance to amikacin, vancomycin, cefepime and moxifloxacin was recorded. In Gram-negative bacteria, a high resistance to 3rd g Cephalosporin's (68.5%) was recorded, while a moderate sensitivity to the other antibiotics tested was recorded.

Conclusion: There is a high rate of undiagnosed LRTI in Yemen and this highlights the need for health authorities to develop strategies to diagnose most of the causes of LRTI, including *Mycoplasma*, *Chlamydia*, and viral causes. No antibiotics are completely effective in treating LRTI in our area and antibiotic sensitivity should be performed in all cases.

Keywords: Antibiotics, Dhamar city, Lower Respiratory Tract Infections (LRTIs), Sana'a City, Yemen.

INTRODUCTION

Lower respiratory tract infection (LRTIs) is the leading cause of infectious diseases of death worldwide, the fifth on the whole cause of death, and the second general reason of disability adjusted life years (DALYs), although they are largely preventable causes

of diseases and Death¹. There have been alterations in the epidemiology of LRTIs in the previous ten years as there has been a reduction in the number of cases among children under 5 and an increase in infection among older adults as well as an increase in viral infections¹. Nevertheless, there is no standardized classification of "LRTIs", a fact which has been said to

impede the admiration of its true epidemiological importance^{2,3}. From an epidemiological standpoint, most definitions of LRTI include influenza, pneumonia, bronchitis (including acute exacerbations of chronic obstructive pulmonary disease [COPD] [AECOPD]) and bronchiolitis as important diseases¹⁻³. The three most important bacterial respiratory pathogens are *Haemophilus influenzae*, *Streptococcus pneumoniae*, and *Moraxella catarrhalis*. Unfortunately, these causes are spreading and increasing the rate of their resistance to antibiotics worldwide^{4,5,6}. The consequence of monitoring the development of this resistance has led to many national, regional and international monitoring programs. Nevertheless, the results of surveillance studies show wide differences in sensitivity rates, both geographically and over time^{7,8}. Bacterial resistance patterns for antibiotics may differ from one region to another depending on the pressure on the antibiotics in that region⁹. Consequently, there is a great need for local resistance spread data in order to guide the experimental prescription and identify areas where new antibiotics with greater effect are needed. In Yemen, data on epidemiology of LRTIs and antibiotic patterns are still rare for bacterial causes. Over the past four years, an increase in mortality has been observed among residents of the capital, Sana'a, due to LRTIs¹⁰. Hence, the current study was planned to isolate the bacterial profile of LRTIs in Yemen and to verify the antibiotic susceptibility among these pathogens in our areas.

SUBJECTS AND METHODS

The selected cases were defined as all patients who had a major complaint of LRTIs and entered the selected Hospitals. The technique of sampling in the study was case finding. As for determining the size of the sample, it was relied on taking all patients who attended selected hospitals during the study period. This study was conducted on 555 hospitalized patients with LRTI in university hospitals in the cities of Sana'a and Dhamar during the period from October 2015 to

October 2018. All patients were subjected to full clinical, radiological and relevant laboratory examinations. Clinical sample analyzes were performed in the laboratories of the National Center of Public Health laboratories Sana'a (NCPHL). The study included 187 patients with community-acquired pneumonia (CAP), 171 patient with hospital-acquired pneumonia (HAP) and 197 patients with acute exacerbation of chronic obstructive pulmonary disease (AECOPD)⁵. CAP was defined as acquired pneumonia outside the hospital¹¹. HAP was defined as a pneumonia occurring 48 hours or more after admission, which was not developed at the time of admission¹². AECOPD were defined according to the GOLD guidelines⁵. Patient data were collected using questionnaire including personal data, clinical symptoms, signs, and history of preexisting chronic diseases. Samples included sputum or bronchoalveolar lavage (BAL) for staining and culture. Samples were cultured on 3 bacteriological media. Blood agar aerobically, chocolate agar with CO₂ and LJ slope then cultures were examined for possible bacterial pathogens of LRTI. Possible bacterial pathogens were isolated and identified using standard laboratory techniques, and microbial sensitivity testing was carried out by means of disc diffusion for selected antibiotics.

Data analysis

The data was statistically analyzed using EPI-Info version 6. The difference in the distribution of bacterial causes among groups was based on a comparison of frequency distributions by chi-square test. The value of $p < 0.05$ was considered significant.

RESULTS

A total of 555 LRTIs hospitalized patients (328/59.1% male and 227/40.9% female) were enrolled in this study. The most frequent age groups were 30-42 years (26.5%), and age group 43-56 years (22.5%); while children age group was less frequent (8.5%).

Table 1: Distribution of age groups, gender and years among LRTI patients and its correlation with bacterial growth outcome.

	NO	% (Total)	Bacterial growth outcome				X ²	P
			Growth		No growth			
			No	%	No	%		
Age category							4.913	0.0296
(3-16)	47	8.5	26	55.3	21	44.7		
(17-29)	119	21.4	76	63.8	43	36.2		
(30-42)	147	26.5	92	62.6	55	37.4		
(43-56)	125	22.5	85	68	40	32		
≥ 57	117	21.1	75	64.1	42	35.9		
Gender							4.940	0.029
Male	328	59.1	218	66.5	110	33.5		
Female	227	40.9	136	60	86	40		
Data							19.124	<0.0001
2015	142	25.6	90	63.4	52	35.6		
2016	178	32.1	124	69.6	54	30.4		
2017	159	28.6	100	62.9	59	37.1		
2018	76	13.7	40	52.6	36	47.4		
Total	555	100.0	354	63.8	201	36.2		

Bacterial growth yielded on 354 (63.8%) while 201 (36.2%) were negative for bacterial culture (Table 1). A large number of LRTI (36.2%) was not diagnosed, mostly in CAP (52.4%), followed by HAP (33.9%) while lower cases were in AECOPD (22.8%). The isolates in 187 patients with CAP were *K. pneumoniae* (26.2%), *S. pyogenes* (12.3%), and *S. pneumoniae* (9%). Isolates in 171 patients with HAP were MSSA (24%), *E. Coli* (12.9%), MRAS (11.1%), *K. pneumoniae* (10.5%) and *P. aeruginosa* (7%). The organisms in 197 patients with AECOPD were *Moraxella catarrhalis* (47.2%), *K. pneumoniae* (17.2%), *H. influenzae* (10.7%) and *P. aeruginosa* (2%) (Table 2). Table 3 shows the frequency of bacterial causative agents of LRTI; the Subtotal Gram positive bacteria were counted for 28.3% from total bacteria isolates, while subtotal Gram positive bacteria was counted for 71.7% from the total bacterial isolates. The most 3 predominant bacteria isolated from LRTIs patients in the study were *K.*

pneumoniae 101(18%), *Moraxella catarrhalis* 91(16.8%) and *S. aureus* 60 (10.8%), while others bacteria such as *S. pyogenes*, *S. pneumoniae*, *H. influenzae*, *P. aeruginosa*, *E. coli* and *Proteus vulgaris* were less frequent (Table 3). In Gram-positive bacteria high resistance was recorded for ampicillin/sulbactam (100%) and amoxicillin/clavulanate (100%), while a moderate sensitivity rate for amikacin, vancomycin, cefepime and moxifloxacin was recorded. In Gram-negative bacteria, a high resistance to 3g of cephalosporins (68.5%) was recorded, while moderate sensitivity to other tested antibiotics was recorded (Table 4). The rates of cure, ICU admission, isolation and death among LRTI cases of positive bacterial growth were almost similar to those of negative culture with slight differences. The mortality rate among total LRTIs was 25%, while for confirmed LRTI cases in bacterial culture it was 22.9%, which is lower among the LRTI cases of negative culture (28.9%) (Table 5).

Table 2: Bacterial profile of lower respiratory tract infections in Yemen.

Common Bacterial pathogens (No/%)		
CAP (n=187/33.7%)	HAP (n=171/30.8%)	AECOPD (n=197/35.5)
<i>S. pneumoniae</i> (17 /9%)	MRSA (19/11.1 %)	<i>H. influenzae</i> (21/10.7%)
<i>K. pneumoniae</i> (49 /26.2%)	<i>k. pneumoniae</i> (18/10.5%)	<i>K. pneumoniae</i> (34/17.2%)
<i>St. pyogenes</i> (23/12.3%)	<i>E. Coli</i> (22/12.9%)	<i>M. catarrhalis</i> (93/47.2%)
	<i>P. aeruginosa</i> (12/7%)	<i>P. aeruginosa</i> (4/2%)
	MSSA (41/24%)	
	<i>Proteus vulgaris</i> (1/0.6%)	
No Bacterial growth (98/52.4%)	No Bacterial growth (58/33.9%)	No Bacterial growth (45/22.8%)

CAP: Community-acquired pneumonia; HAP: Hospital-acquired pneumonia; AECOPD: Acute exacerbations of chronic obstructive pulmonary disease; MRSA: Methicillin-resistant *Staphylococcus aureus*; MSSA: Methicillin-sensitive *Staphylococcus aureus*.

DISCUSSION

Lower respiratory tract infection (LRTIs) is the leading cause of infectious diseases of death worldwide, the fifth general cause of death, and the second general cause of disability adjusted life years (DALYs), although they are largely preventable causes of diseases and Death¹. In the current study the mortality rate among total LRTIs was 25%, while for LRTIs cases confirmed for bacterial culture was 22.9%, lower than that among negative culture LRTI cases (28.9%) (Table 5); this rate is higher than that reported by Brown and others in the United States of America where the death rate among community-acquired pneumonia hospitalizations patients was 7.4%¹³. While Global Strategy for the Diagnosis, Management and Prevention of COPD, reported that long-term prognosis following LRTIs was poor, with a 5-year mortality of approximately 50%¹⁴. The high mortality rate in the current study may be high rates for related factors and include comorbidities especially cardiovascular disease, severity of exacerbations¹. When reviewing the various studies, it is clear that there are some regional

differences in the reported etiology of LRTIs, as described by Waterer¹⁵.

Table 3: The frequency of bacterial causative agents of LRTI.

Isolated Bacteria	No (%)	% Total n=555
Gram Positive		
<i>S. pneumoniae</i>	17 (17)	3
<i>S. aureus</i>	60 (60)	10.8
<i>S. pyogenes</i>	23 (23)	4.1
Subtotal Gram positive	100 (28.3)	18
Gram Negative		
<i>K. pneumoniae</i>	101 (39.8)	18
<i>H. influenzae</i>	21 (8.3)	3.8
<i>P. aeruginosa</i>	16 (6.3)	2.9
<i>Proteus vulgaris</i>	1 (0.4)	0.18
<i>Moraxella catarrhalis</i>	93 (36.6)	16.8
<i>E. coli</i>	22 (8.7)	4
Subtotal Gram negative	254 (71.7)	45.8
Total positive culture	354	63.8
Total negative culture	201	36.2
Fungi		
<i>C. albicans</i> colonization	159	28.6

This may be related to a number of factors, but it is also important to realize that although LRTIs are not a seasonal disease, many different organisms, including *S. pneumoniae*, influenza virus, do have seasonal variations¹⁶. In the current study the most 3 predominant bacteria isolated from LRTIs patients were *K. pneumoniae* 101(18%), *Moraxella catarrhalis* 91(16.8%) and *S. aureus* 60(10.8%), while *S. pneumoniae* and *H. influenzae* were less frequently (Table 3); this result is different from that traditionally, the *S. pneumoniae* has been reported to be the mainly widespread cause of LRTIs¹⁷⁻¹⁹ and the Global Burden of Disease Study analysis of LRTIs

(2015)¹ revealed that the *S. pneumoniae* was the most common cause of LRTIs among all ages. In spite of this, current results matches with reports in which there have been changes noted in the reported etiology of LRTIs, particularly with the use of more sensitive diagnostic tools¹⁹⁻²¹. In wide-ranging, it is gradually more recognized that viruses look to play a bigger role in the etiology of LRTIs than has previously been documented²²⁻²⁵ and cases of infection with more than one pathogen, commonly the association of one or more viruses with one or more bacterial agents are not uncommon^{21,22}.

Table 4: Antibiotic sensitivity and resistance rates (percentages) of gram positive and gram negative bacteria in 354 patients with LRTI in Yemen.

Antibiotics	Test	Bacteria	
		Gram positive percentage	Gram negative Percentage
Vancomycin	S	69.2	ND
	I	11	
	R	19.8	
Moxifloxacin	S	47.5	71
	I	14	7
	R	38.5	22
Cephalosporin's	S	8	22.5
	I	13	9
	R	79	68.5
Ciprofloxacin	S	37	82
	I	12	3.2
	R	51	14.8
Cefepime	S	46	65.4
	I	19	13.1
	R	35	21.5
* Aampicillin/ sulbactam	S	0	48.2
	I	0	11.2
	R	100	40.6
*Amoxicillin/ clavulanate	S	0	67
	I	0	11
	R	100	22
Amikacin	S	58.2	80
	I	19	9
	R	22.8	11

*Not done for *P. aeruginosa*, ND= not done

Table 5: The output of LRTI cases with bacterial infections in comparison with LRTI cases caused by other agents.

Outcome	LRTI with bacterial infections (n=354)		LRTI cases with non-bacterial agents (n=201)		Total (n=555)	
	No	%	No	%	No	%
Cure	273	77.1	143	71.1	416	75
ICU	22	6.2	31	15.4	53	9.5
Isolation	1	0.3	3	1.5	4	0.72
Death	81	22.9	58	28.9	139	25
Total n=555	354	63.8	201	36.2	555	100

For patients with CAP, current results (Table 2) showed bacterial profiles similar to those reported by international studies⁶ and regional²⁶. This pattern of "local" hegemony should be taken into account when prescribing antimicrobials in our region. When antibiotic sensitivity was considering for bacterial isolates from LRTI patients, in Gram positive bacteria a high resistance was recorded for ampicillin/sulbactam (100%) and amoxicillin/clavulanate (100%), while

moderate of sensitivity was recorded for amikacin, vancomycin, Cefepime and moxifloxacin. In Gram negative bacteria a high resistance was recorded for 3rd g Cephalosporin's (68.5%), while moderate of sensitivity was recorded for other tested antibiotics (Table 4). Current data revealed high resistance rates for cephalosporins, and the β -lactam- β -lactamase inhibitors. These findings are in agreement with the increasing prevalence of resistance of Gram positive

bacteria as *S. pneumoniae* to those antimicrobial groups, by regional^{7,27,29}, and worldwide^{6,7} studies. Moreover, current results highlight the increasing problem of MDR in Gram positive and Gram negative bacteria of LRTIs, a problem that was extensively addressed in the literature²⁸⁻³⁰. This warns us of the need for wise use of different groups of antimicrobials, especially in our resource-poor country. Moreover, this requires greater focus on identifying drivers of resistance relevant and on implementing effective strategies to combat resistance and MDR problems. For patients with HAP, the difficulty of antibiotic resistance appears more important; as a result, the situation is more complex than that in CAP. Nosocomial pneumonias leads to high morbidity and mortality, in particular amongst ICU patients^{8,11}. In most clinical cases, it is needed to start empirical antimicrobial therapy before obtaining microbial results. On the other hand, the situation is further complicated by the emergence of several beta-lactamase and MDR pathogens^{29,31}. Obviously there is a great need to obtain data on the prevalent strains in HAP; along with the sensitivity pattern to help revise antibiotic policy and guide physicians to better manage patients with HAP; especially in developing countries such as Yemen. The current study revealed the present of MRSA, Gram-negative organisms, and *P. aeruginosa* among patients with HAP. This differs clearly from the results obtained by Goel and co-workers³¹ and even those of Ahmed, *et al.*,³², Agmy, *et al.*,³³. Although the later study addressed the problem of HAP in 75 cases of ICU patients, the predominant pathogens were *S. aureus* (32%), *P. aeruginosa* (30%), and *S. pneumoniae* (15%). It is clear that this "regional" difference explains the changing pattern of pathogens that cause over time, even in the same hospital. This underscores the importance of implementing continued local monitoring programs⁸. Also, current data show an alarming high prevalence of MRSA. This coincides with the recent report by Alyahawi, and Al-Safani *et al.*,^{34,35} who observed that the prevalence of MRSA in invasive isolates from hospitals in Yemen was 23%³⁴.

CONCLUSIONS

Lower respiratory infections are still very common and continue to be a major cause of morbidity and mortality in Yemen in children and adults alike, and there are significant changes in the epidemiology of LRTIs in terms of their frequency and infectious pathogens. There is a high rate of undiagnosed LRTI in Yemen and this highlights the need for health authorities to develop strategies to diagnose most of the causes of LRTI, including *Mycoplasma*, *Chlamydia*, and viral causes. The most common bacteria in CAP in Yemen is *K. pneumoniae* while HAP is the *S. aureus* and Gram negative bacteria. For acute exacerbation of COPD, *M. catarrhalis* was the most common. No antibiotics are completely effective in treating LRTI in our area and antibiotic sensitivity should be performed in all cases.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Ministry of Health and Population, Sana'a, Yemen and the National Center of Public Health Laboratories (NCPHL), Ministry of Health and Population, Sana'a, Yemen for their support and provided working space and materials.

AUTHOR'S CONTRIBUTION

This research work is part of project of the National Center for Public Health Laboratories (NCPHL). The authors performed clinical and laboratory works. **Al-Haifi YA:** writing original draft, methodology. **Al Makdad ASM:** investigation, formal analysis, conceptualization. **Salah MK:** editing. **Al-Shamahy HA:** supervision, conceptualization. **Al Shehari WAA:** writing, review, and editing, methodology. **Gunduz C:** writing, review, and editing. All authors revised the article and approved the final version.

DATA AVAILABILITY

The data supporting the findings of this study are not currently available in a public repository but can be made available upon request to the corresponding author.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. GBD 2015 LRI Collaborators. Estimates of the global, regional and national morbidity, mortality, and aetiologies of lower respiratory tract infections in 195 countries: a systematic analysis for the global burden of disease study 2015. *Lancet Infect Dis* 2017; 17:1133–1161.
2. ERS white book, Chapter 18. Acute lower respiratory infections. ERS white book. p. 210–223. [cited 2020 April 21]. Available from: https://www.erswhitebook.org/files/public/Chapters/18_ALRIs.pdf
3. Greene G, Hood K, Little P, *et al.* Towards clinical definitions of lower respiratory tract infection (LRTI) for research and primary care practice in Europe: an international consensus study. *Prim Care Respir J* 2011; 20(3):299–306. <https://doi.org/10.4104/pcrj.2011.00034>
4. Pfaller MA, Jones RN, Doren GV, *et al.* Bacterial pathogens isolated from patients with bloodstream infection: frequencies of occurrence and antimicrobial susceptibility patterns from the SENTRY antimicrobial surveillance program (United States and Canada, 1997). *Antimicrob Agents Chemother* 1998; 42: 1762-1770. PMID:9661018 PMID:PMC105680
5. Global Initiative for Chronic Obstructive Lung Disease. Global Strategy For The Diagnosis And Prevention Of Chronic Obstructive Pulmonary Disease. Updated 2009.
6. Mandell LA, Wunderink RG, Anzueto A, *et al.* Infectious diseases society of america/american thoracic society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis* 2007; 44 Suppl 2: S27–72. <http://dx.doi.org/10.1086/511159> PMID:17278083
7. Garcí'a-Rey C, Aguilar L, Baquero F, *et al.* Importance of Local Variations in Antibiotic Consumption and Geographical Differences of Erythromycin and Penicillin

- Resistance in *Streptococcus pneumoniae*. J Clin Microbiol 2002; 40: 159–164.
<http://dx.doi.org/10.1128/JCM.40.1.159-164.2002>
8. Alpuche C, Garau J, Lim V. Global and local variations in antimicrobial susceptibilities and resistance development in the major respiratory pathogens. Int J Antimicrob Agents 2007; 30 Suppl 2: 135-8.
<http://dx.doi.org/10.1016/j.ijantimicag.2007.07.035>
 9. Lakshmi V. Need for national/regional guidelines and policies in India to combat antibiotic resistance. Indian J of Medical Microbiology 2008; 26: 105-7.
<http://dx.doi.org/10.4103/0255-0857.40521>
 10. Al-Ademi DAA, Al-Harazi AH, Al-Shamahy HA, Jaadan BM. Detection of influenza viruses among hospitalized cases suffering from severe acute respiratory illness (SARI) in Sana'a city, Yemen. Universal J Pharm Res 2019; 4(4): 30-34. <https://doi.org/10.22270/ujpr.v4i4.296>
 11. Gordon RC. Community acquired pneumonia in adolescents. Adoles Med 2000; 11: 681-695. PMID:11060562.
 12. American Thoracic Society; Infectious Diseases Society of America. Guidelines for the Management of Adults with Hospital acquired, Ventilator- associated, and Healthcare-associated Pneumonia. Am J Res Crit Care Med 2005; 171: 388–416. <http://dx.doi.org/10.1164/rccm.200405-644ST>
 13. Brown JD, Harnett J, Chambers R, et al. The relative burden of community-acquired pneumonia hospitalizations in older adults: a retrospective observational study in the United States. BMC Geriatr 2018; 18(1):92. <https://doi.org/10.1186/s12877-018-0787-2>
 14. GOLD 2017: Global Strategy for the Diagnosis, Management and Prevention of COPD, Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017. Available from: <https://goldcopd.org>.
 15. Waterer G. Community-acquired pneumonia: a global perspective. Semin Respir Crit Care Med 2016; 37:799–805. <https://doi.org/10.1055/s-0036-1592313>
 16. Feldman C, Anderson R. The role of *Streptococcus pneumoniae* in community-acquired pneumonia. Semin Respir Crit Care Med 2016; 37:806–818. <https://doi.org/10.1055/s-0036-1592074>
 17. Welte T, Torres A, Nathwani D. Clinical and economic burden of community-acquired pneumonia among adults in Europe. Thorax 2012; 67:71–79. <http://dx.doi.org/10.1136/thx.2009.129502>
 18. Herrero FS, Olivás JB. Microbiology and risk-factors for community-acquired pneumonia. Semin Respir Crit Care Med 2012; 33:220–231. <http://dx.doi.org/10.1055/s-0032-1315634>
 19. Cillóniz C, Cardozo C, García-Vidal C. Epidemiology, pathophysiology, and microbiology of community-acquired pneumonia. Ann Res Hosp 2018; 2:1. <http://dx.doi.org/10.21037/arih.2017.12.03>
 20. Lung M, Rello J. Microbiology of bacterial CAP using traditional and molecular techniques. Eur Res Mon 2014; 63:25–41. <http://dx.doi.org/10.1183/1025448x.10003213>
 21. Gadsby NJ, Russell CS, McHugh MP, et al. Comprehensive molecular testing for respiratory pathogens in community-acquired pneumonia. Clin Infect Dis 2016; 62(7):817–823. <http://dx.doi.org/10.1093/cid/civ1214>
 22. Jain S, Self WH, Wunderink RG, et al. Community-acquired pneumonia requiring hospitalization among U.S. Adults N Engl J Med 2015;373(5):415–427. <http://dx.doi.org/10.1056/NEJMoa1500245>
 23. Da Rocha Neto OG, Leite RF, Baldi BG. Update on viral community-acquired pneumonia. Rev Assoc Med Bras 2013; 59 (1):78–84. [https://doi.org/10.1016/S2255-4823\(13\)70433-8](https://doi.org/10.1016/S2255-4823(13)70433-8)
 24. Rohde GGU. The role of viruses in CAP. Eur Respir Monogr 2014; 63:74–87. <https://doi.org/10.1183/1025448x.10003513>
 25. Jain S. Epidemiology of viral pneumonia. Clin Chest Med 2017; 38:1–9. <https://doi.org/10.1016/j.ccm.2016.11.012>
 26. El Sayed Zaki M and Goda T. Clinico-pathological study of atypical pathogens in community-acquired pneumonia: A prospective study. J Infect Dev Ctries 2009; 3: 199-205. <http://dx.doi.org/10.3855/jidc.36>
 27. Agmy G, Mohamed S, Gad Y, et al. Bacterial profile, antibiotic sensitivity and resistance of lower respiratory tract infections in upper Egypt. Mediterr J Hematol Infect Dis 2013, 5(1): e2013056, <http://dx.doi.org/10.4084/MJHID.2013.056>
 28. Wasfy MO, Pimentel G, Abdel-Maksoud M, et al. Antimicrobial susceptibility and serotype distribution of *Streptococcus pneumoniae* causing meningitis in Egypt, 1998–2003. J Antimicrob Chemother 2005; 55: 958–964. <http://dx.doi.org/10.1093/jac/dki101>
 29. Borg MA, Tiemersma E, Scicluna E, et al. Prevalence of penicillin and erythromycin resistance among invasive *Streptococcus pneumoniae* isolates reported by laboratories in the southern and eastern Mediterranean region. Clin Microbiol Infect 2009; 15: 232–237. <http://dx.doi.org/10.1111/j.1469-0691.2008.02651.x>
 30. Lynch JP 3rd, Zhanel GG. *Streptococcus pneumoniae*: Does antimicrobial resistance matter? Semin Res Crit Care Med 2009; 30: 210-38. <http://dx.doi.org/10.1055/s-0029-1202939>
 31. Goel N, Chaudhary U, Aggarwal R, Bala K. Antibiotic sensitivity pattern of gram negative bacilli isolated from the lower respiratory tract of ventilated patients in the intensive care unit. Indian J Crit Care Med 2009; 13: 148-151. <http://dx.doi.org/10.4103/0972-5229.58540>
 32. Ahmed SH, Daef EA, Badary MS, et al. Nosocomial blood stream infection in intensive care units at Assiut University Hospitals (Upper Egypt) with special reference to extended spectrum beta-lactamase producing organisms. BMC Research Notes 2009; 2: 76-87. <http://dx.doi.org/10.1186/1756-0500-2-76>
 33. Agmy GM, Sayed SS, Mohamed EA. Nosocomial pneumonia in intensive care units at Assiut University Hospital. In Proceedings of the Annual Congress of The European Respiratory Society, Copenhagen; 2005. PMID:19498617
 34. Al-Safani AMA, Al-Shamahy HA, Al-Moyed KA. Prevalence, antimicrobial susceptibility pattern and risk factors of MRSA isolated from clinical specimens among military patients at 48 medical compound in Sana'a city-Yemen. Universal J Pharm Res 2018; 3(3): 40-44. <https://doi.org/10.22270/ujpr.v3i3.165>
 35. Alyahawi A, Alkaf A, Alhomidi A. Prevalence of methicillin resistant *Staphylococcus aureus* (MRSA) and antimicrobial susceptibility patterns at a private hospital in Sana'a, Yemen. Universal J Pharm Res 2018; 3(3): 4-9. <https://doi.org/10.22270/ujpr.v3i3.159>