



RESEARCH ARTICLE

COMPARISON BETWEEN THE EFFECTS OF THYMOQUINONE OBTAINED FROM THE SEEDS OF *NIGELLA SATIVA* AND VERAPAMIL ON THE VOLUME AND ACIDITY OF GASTRIC ACID SECRETION

Rahma Hamayun¹, Muhammad Jan¹, Muhammad Fazeel²

¹College of Medicine, Northern Border University, Arar, Kingdom of Saudi Arabia.

²University of Agriculture, Pakistan.

Article Info:



Article History:

Received: 9 June 2020
 Reviewed: 13 July 2020
 Accepted: 24 August 2020
 Published: 15 September 2020

Cite this article

Hamayun R, Jan M, Fazeel M. Comparison between the effects of thymoquinone obtained from the seeds of *Nigella sativa* and verapamil on the volume and acidity of gastric acid secretion. Universal Journal of Pharmaceutical Research 2020; 5(4):57-60.

<https://doi.org/10.22270/ujpr.v5i4.453>

*Address for Correspondence:

Dr. Rahma Hamayun, College of Medicine, Northern Border University, Arar, Kingdom of Saudi Arabia, Tel: +966547282585, E-mail: r_hamayun@hotmail.com

Abstract

Background and Objectives: The over production of gastric acid results in peptic ulcer. This study was done to compare the effects of thymoquinone and Verapamil on volume and acidity of carbachol induced gastric secretion.

Methods: There were 24 rabbits used, weighing 1-1.5 kg. The rabbits were kept on fasting for 48 hours. After fasting, the pylorus of each rabbit was ligated. Thymoquinone 5 mg/kg, Carbachol 600 µg/kg and Verapamil 10 mg/kg body weight were administered intraperitoneally. Pylorus ligation method was used for getting gastric contents and titration method was used for finding out acidity.

Results: Verapamil has been proved very effective for the treatment of many diseases. The drug verapamil inhibits the release of histamine, acetylcholine and gastrin. Verapamil has also shown effects in reducing the secretion of gastric acid. It was found that Thymoquinone reduced the volume, free and total acidity of gastric secretion, which were statistically highly significant when compared with Carbachol ($p=0.000$) but when we compared the results of Thymoquinones with that of Verapamil, it was non-significant.

Conclusions: It was concluded that Thymoquinone can be used for the treatment of peptic ulcer and all other diseases which are caused by increased gastric acidity like dyspepsia, gastritis and reflux esophagitis.

Keywords: Gastric secretion, Thymoquinone, Verapamil.

INTRODUCTION

In clinical practice, peptic ulcer is one of the most common medical complain. In majority of the patients, peptic ulcer is caused by elevated acid production from gastric mucosa. In patients who are achlorhydric, ulcers are not found. Ulcers mostly occur in Zollinger-Ellison (Z.E) syndrome which is caused by excess acid secretion¹. The goal of treatment of peptic ulcer is to inhibit the over production of acid. *Nigella sativa* is a part of the botanical family of Ranunculaceae. It is commonly cultivated in Europe, Middle East and Westren Asia. All over the world, it is called by many names like habbat al-baraka or Kali jeera. In the light of Hadith "Use this Black seed, it has a cure for every disease except death" (Sahih Bukhar), The *Nigella sativa* (*N. sativa*) seeds, are mostly used in many Arab countries like Saudi Arabia, Middle Eastas a natural remedy for many diseases. *N. sativa* seeds contain many active ingredients, the most important of which is thymoquinone (nigellone)². Because of the large

variety of uses of *N. sativa*, many researchers have conducted various *in vitro* and *in vivo* studies on laboratory animals and human beings in order to find out their pharmacological activities. These include anti-inflammatory³, analgesic, anti-pyretic activity⁴, antimicrobial^{5,6,7,8}, antifungal^{9,10,11}, hypoglycemic¹² and anti-tuberculosis¹³.

Thymoqui-nione administration can prevent and improve the murine-DSS (dextran sodium sulfate) induced colitis. It could also serve as an effective therapeutic agent for the treatment of patients with inflammatory bowel disease. It also helps to prevent colitis and diarrhea in patients¹⁴.

MATERIALS AND METHODS

The active ingredient thymoquinone was obtained from Amidis chemical company PVT limited, China. Thymoquinone was extracted from the *N. sativa* plant by the company itself.

Source of chemical:

The chemical verapamil was purchased from VPL chemicals, private limited India.

Method

Twenty-four rabbits of local breed were selected for this study. The animals were healthy and were of both sexes. All the chemicals were administered through intraperitoneal route according to the body weight of the animals. The animals were not given any food for 48 hours. The animals were separated into 3 groups each group had 8 animals. Group 1 was administered Carbachol. The dose of carbachol was 600 µg/kg body weight, Group 2 was administered Thymoquinone with a dose of 5 mg/kg and group 3 was given Verapamil. The dose of verapamil was 10 mg/kg body weight. After 15 minutes Carbachol with a dose of 600 µg/kg body weight was administered to Group 2 and 3. The gastric juice was obtained from all rabbits by the method of pylorus ligation, as explained by Vischer et al.,¹⁵. The anesthesia was given with ether in a big glass desicator, and all the animals were weighted. The abdomen was cut by an incision in mid-line and the pylorus was disconnected with silk suture. Suture clamps were used to close the wall of abdomen. The inhibitory effect of the drug was better understood after the stimulation produced by carbachol. When the anesthesia was stopped, the animals became conscious. After a period of 4 hours, all the animals were slaughtered, their abdomen was again opened and the cardiac end of the stomach was ligated. The incision

was given to stomach at greater curvature. The gastric juice was finally procured and was titrated against 0.1 N NaOH solution by the procedure explained by Varley¹⁶. This procedure is being performed since 1954, for the calculation of all forms of acidity i.e. free, combined and total. In this process, one ml of centrifuged gastric juice is titrated against 0.1 N NaOH using Topfer's reagent as an indicator for calculation of free acidity and 1% phenolphthalein as a standard for combined acidity. Total acidity was calculated by the sum of the two titrations.

Statistical analysis

The data obtained was subjected to statistical analysis for any significance. The data was entered into SPSS-IBM Version 19. *p* value of <0.05 was considered to be statistically significant.

RESULTS

The volume, free acidity and total acidity in group 1 were 28.125±2.031 ml, 6.225±1.188 m.Eq./dl and 7.650±1.243 m.Eq./dl respectively. Similarly, the mean values for volume, free acidity and total acidity of gastric secretion in group 2 (Thymoquinone+ Carbachol treated group) were 13.625±1.355 ml, 2.412±.626 m.Eq./dl and 3.750±.833 m. Eq./dl respectively. There was a reduction in all the parameters and was found to be highly significant when compared with Carbachol group (*p*=0.000). All these changes are shown in Table 1.

Table 1: Effects of Thymoquinone and Verapamil on the volume and acidity of gastric secretion induced by Carbachol in fasting rabbits. (*p*=0.000)

Drugs	Volume of gastric secretion (ml)	Acidity (meq/dl of gastric secretion)	
		Free	Total
Carbachol	28.125±2.031	6.225±1.188	7.650±1.243
Thymoquinone +Carbachol	13.625±1.355	2.412±.626	3.750±.8
<i>p</i> Values (When compared with carbachol)	0.000	0.000	0.000
Verapamil+ Carbachol	13.212±1.501	2.200±.575	3.575±.497
<i>p</i> Values (When compared with Carbachol)	0.000	0.000	0.000

Each value indicates mean of the total observation. *p* Value between Carbachol and drugs+ Carbachol

Likewise, the mean values for volume, free acidity and total acidity of gastric secretion in group 3 (Verapamil + Carbachol treated group) were 13.212±1.501 ml, 2.200±.575 m.Eq./dl and 3.575±.497 m.Eq./dl respectively. There was a reduction in all the parameters and was found to be highly significant when compared with Carbachol group (*p*=0.000). All these changes are shown in Table 1. Similarly, when we compared the mean values for volume, free and total acidity of Verapamil+ Carbachol group to those of Thymoquinone + Carbachol group, *p* values were 0.392, 0.204, 0.412 (Table 2).

DISCUSSION

N. sativa seed and its constituents are mostly utilized as a natural cure for many diseases. A lot of scientific work has been conducted to determine the pharmacological activities of this plant. Most of the

research has established its importance in traditional medicine as an analgesic, anti-inflammatory, anti-oxidant, anti-cancer, anti-microbial, anti-parasitic, antihypertensive and as an immune booster. The main neurotransmitters/hormones that directly increase the secretion by the gastric organs are acetylcholine, gastrin and histamine¹⁷. The stimulation of these neurotransmitters depends on the influx of Ca ions¹⁸. The increase in gastric volume and acidity is due to the intravenous administration of calcium which results in hypercalcemia¹⁹. During an *in vitro* study, it was explained that *Nigella sativa*, successfully stopped the release of histamine from mast cells, through lowering the intracellular calcium and blockage of protein kinase C. In a study relating to hypertensive rats, it was found that *N. sativa* extract produced a significant hypotensive effect when compared to that of 0.5 mg/kg/day of oral calcium channel blocker nifedipine²⁰.

Table 2: Comparison between the effect of Thymoquinone and Verapamil on the volume and acidity of gastric secretion induced by Carbachol in fasting rabbits.

Drugs	Volume of gastric secretion (ml)	Acidity (m.Eq/dl of gastric secretion)	
		Free	Total
Thymoquinone+Carbachol	13.625±1.355	2.412±.626	3.750±.833
Verapamil+Carbachol	13.212±1.501	2.200±.575	3.575±.497
P values	0.392	0.204	0.412

Each value indicates mean of the total observation. *p* Value between Thymoquinone+Carbachol and Verapamil+Carbachol

N. sativa antagonized methacholine induced contractions of isolated guinea-pig tracheal chain²¹. This study explained the anticholinergic effect of *N. sativa* which could be the reason for inhibiting the gastric acid secretion. Further, it was found that this study is reported to show significant response as concluded by other scientists who found that calcium channel blocker Verapamil significantly reduces gastric acid secretion^{22,23}. Calcium channel blockers block the calcium influx, which causes a reduction in volume and acidity of gastric juices. Also, the lipoxygenase pathway, a step-in metabolism of arachidonic acid, is also blocked by calcium channel blockers. As a result, the leukotrienes, the harmful substance is not formed and all the arachidonic acid is metabolized by cyclooxygenase pathway. This cause the production of prostaglandin which couples with guanine nucleotide binding protein (Gi protein) and blocks the adenyl cyclase and thus decreases the gastric secretion²⁴. Release of histamine from mast cells is critically dependent on external calcium ions, so by blocking calcium ions can block the release of histamine. Histamine is an important factor for increasing the gastric acid secretion²³. In this study it was investigated that thymoquinone, which is obtained from *N. sativa* showed maximal reduction in gastric secretion and acidity. This study can be correlated to the study done by El-Dakhkhani *et al.*,²⁵. El-Dakhkhani *et al.*, observed the effect of *N. sativa* oil on HCL secretion and ethanol-induced ulcer in rats. The gastroprotective and anti-secretory effects of *N. sativa* seed powder significantly reduced gastric secretion, volume, pH and gastric acid-output. This was explained as the acid reducing effect of thymoquinone is due to its antihistaminic effect. Histamine is the main stimulus for histamine release that cause high gastric hydrochloric acid secretion leading to peptic ulcer, gastritis etc. Thymoquinone inhibits histamine, also increase gastric mucous secretion which is protective to stomach²⁶. Significant increase in mucin content, glutathione level as well as a significant decrease in mucosal histamine content and ulcer formation. From above discussion it is clear that thymoquinone obtained from the seeds of *N. sativa* can significantly reduce the gastric acidity by blocking the release of histamine from histamine H2 receptors. It also contains calcium channel blocking activity which reduces the release of acetylcholine and histamine.

CONCLUSIONS

It is concluded that thymoquinone obtained from the *N. sativa* seeds may be effectively used in patients with diseases of peptic ulcer and other medical conditions

caused by excess secretion of HCL. For the evaluation of these effects, further experiments should be done in human subjects. Current study concluded that thymoquinone significantly decreased the Carbachol stimulated acid secretion. Further work on the mechanism of action is suggested.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Northern Border University, Arar, Kingdom of Saudi Arabia to provide necessary facilities for this work.

AUTHOR'S CONTRIBUTION

Hamayun R: writing original draft, methodology. **Jan M:** investigation, conceptualization. **Fazeel M:** formal analysis, interpretation of data. All authors revised the article and approved the final version.

DATA AVAILABILITY

The datasets generated during this study are available from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. Edward, CRW, Bouchier, IAD and Haslett, C. Diseases of the stomach. In: Davidson's Principles and Practice of Medicine, Churchill Livingstone, London, 1995: 425-434.
2. El-Zawahry BH: Chemical composition of *Nigella sativa* Linn seed. In: effect of *Nigella sativa* on certain aspects of metabolism after feeding normal and hyperlipidemic diet in adult and old animals [MD thesis in medical basic sciences]. Cairo, Egypt: Al-Azhar University, 1997:16-28.
3. Houghton PJ, Zarka R, de las Heras B, Hoult JR. Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane lipid peroxidation. *Planta Med* 1995; 61(1): 33-36. <https://doi.org/10.1055/s-2006-957994>
4. Al-Ghamdi MS. Anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. *J Ethnopharmacol*. 2001; 76: 45-8. [https://doi.org/10.1016/s0378-8741\(01\)00216-1](https://doi.org/10.1016/s0378-8741(01)00216-1)
5. Topozada HH, Masloum H and El-Dakhkhany M. The anti-bacterial properties of *Nigella sativa* seeds. Active principle with some clinical application. *J Egypt Med Assoc* 1995; 48: 187-202. *PMID: 5873673*

6. El-Fatraty. Isolation and structure assignment of an antimicrobial principle from the volatile oil of *Nigella sativa* L seeds. *Pharmazie* 1975; 30(2): 109-11. *PMID: 238225*
7. Roy J, Shakleya D, Callery PS, Thomas JG. Chemical constituents and antimicrobial activity of a traditional herbal medicine containing garlic and black cumin. *African J Trad, Comp Alt Med* 2006; 3 (2): 1-7.
8. Randhawa M. An update on antimicrobial effect of *Nigella sativa* and experience at King Faisal University Dammam Saudi Arabia. *J Saudi Soc Dermal Dermatol Surg* 2008; 1237-44.
9. Roy J, Shakleya D and Aljabre S. *In vitro* antifungal activity of thymoquinone against *Scopulariopsis brevicaulis*. *Arab J Pharm Sci* 2005; 3: 27-33.
10. Akhtar N, Aljabre S. The effect of thymoquinone and amphotericin B on the growth of *Aspergillus niger*. *Scientific J King Faisal University* 2007; 8 (1):4.
11. Daba M, Abdel-Rehman M. Hepatoprotective activity of thymoquinone in isolated rat hepatocytes. *Toxicol Lett* 1998; 95 (1): 23-9.
[https://doi.org/10.1016/s0378-4274\(98\)00012-5](https://doi.org/10.1016/s0378-4274(98)00012-5)
12. Al-Awadi FM, Gumma KA. Studies on the activity of individual plants of an anti-diabetic plant mixture. *Acta Diabetol Lat* 1987; 24(1): 37-41.
<https://doi.org/10.1007/BF02732051>
13. Randawa MA. *In vitro* anti tuberculous activity of thymoquinone an active principal of *Nigella sativa*. *J Ayub Med Coll* 2011; 23(2): 78-81. *PMID: 24800349*
14. Xiaofei L, Meng L, Zirong Y, Mengyao J, Xufeng G and Weiguo D. Thymoquinone prevents and ameliorates dextran sulfate sodium-induced colitis in mice. *Digest Dis Sci* 2012; 57 (9): 2296-2303.
<https://doi.org/10.1007/s10620-012-2156-x>
15. Vischer FE, Seay PH, Tazelaar AP, Veldkamp Jr W, Brook MJ. Pharmacology of Pamine Bromide. *J Pharmacol Expt Therapy* 1954; 110: 118-204.
PMID: 13118493
16. Varley H: Test of gastric function, occult blood. In: *Practical clinical Biochemistry*: London, Williams Meinmann 1962: 249-277.
[https://doi.org/10.1016/0307-4412\(82\)90103-0](https://doi.org/10.1016/0307-4412(82)90103-0)
17. Hayward NJ, Harding M, Liyod SAC, *et al.* The effect of CCKB/gastrin antagonists on stimulated gastric acid secretion in the anesthetized rat. *British J Pharmacol* 1991; 104: 973-977.
<https://doi.org/10.1111/j.1476-5381.1991.tb12535.x>
18. Berglindh T, Sachs G, Tacheguchi N. Ca⁺⁺ dependent secretagogue stimulation isolated rabbit gastric glands. *Am J Physiol* 1980; (239): 90-94.
<https://doi.org/10.1152/ajpgi.1980.239.2.G90>
19. Passaro E Jr, Basso N, Walsh JH. Calcium challenge in Zollinger- Ellison syndrome. *Surgery* 1972; 72:60-7.
20. Zaoui A, Cherrah Y, Aloui K, Mahassine N, *et al.* Effect of *Nigella sativa* fixed oil on blood homeostasis in rat. *J Ethnopharmacol* 2002; 79(1): 23-6.
[https://doi.org/10.1016/S0378-8741\(01\)00342-7](https://doi.org/10.1016/S0378-8741(01)00342-7)
21. Boskabady MH, Shahabi M. Bronchodilatory and anticholinergic effects of *Nigella sativa* on isolated guinea pig tracheal chains. *Iran J Med Sci* 1997; 22(3, 4): 133.
22. Rogers C, Pihan, G, Szabo S. Role of leukotrienes in the pathogenesis of haemorrhagic gastric mucosal lesions induced by ethanol or HCl in the rat. *Gastroenterol* 1986; 90:1797. <https://doi.org/10.1007/bf01536761>
23. Main IHM, Pearce JB. Effects of calcium on acid secretion from the rat isolated gastric Mucosa during stimulation with histamine, pentagastrin, methacholine, and dibutyl cyclic adenosine-3,5monophosphate. *Br, J Pharmacol* 1978; 64: 359-368. *PMID: 214194*
24. Flekanstein A. Specific pharmacology of calcium in pericardium, cardiac pacemaker and vascular smooth muscles. *Ann Rev Pharmacol Toxicol* 1977; 17:149-166.
<https://doi.org/10.1146/annurev.pa.17.040177.001053>
25. El-Dakhkhani M, Barakat M, El-Halim MA, Aly SM. Effect of *Nigella sativa* oil on gastric secretion and ethanol-induced ulcer in rats. *J Ethnopharmacol* 2000; 72 (1-2): 299-304.
[https://doi.org/10.1016/s0378-8741\(00\)00235-x](https://doi.org/10.1016/s0378-8741(00)00235-x)
26. Shakeri F, Gholamnezhad Z, Mégarbane B, Rezaee R, Boskabady MH. Gastrointestinal effects of *Nigella sativa* and its main constituent, thymoquinone: a review. *Avicenna J Phytomed* 2016; 6(1): 9–20.
PMID: 27247918